

PUBLIC HEALTH REPORTS

VOL. 46

DECEMBER 25, 1931

NO. 52

TYPHUS FEVER: TYPHUS VIRUS IN FECES OF INFECTED FLEAS (*XENOPSYLLA CHEOPIS*) AND DURATION OF INFECTIVITY OF FLEAS

By E. T. CEDER, Assistant Surgeon, R. E. DYER, Surgeon, and A. RUMREICH, and L. F. BADGER, Passed Assistant Surgeons, U. S. Public Health Service

As a step in the elucidation of the mechanism by which the rat flea (*Xenopsylla cheopis*) transmits endemic typhus fever of the United States from rat to rat, or from rat to man, experiments have been made to determine the presence of the virus in the feces of infected fleas. As noted in a previous publication (1), the feces of fleas infected by feeding on white rats which had been inoculated with the virus of endemic typhus were found to be infectious. The experiments bearing on this point follow:

Rat fleas (*Xenopsylla cheopis*) were placed in one of the glass boxes previously used in transmission experiments (2) (3). White rats were inoculated with the virus of endemic typhus and introduced into the box which contained the fleas. After a period of two weeks a few fleas were removed, ground up in salt solution, and injected into 2 guinea pigs. The reaction typical of endemic typhus resulted in both injected animals. Approximately 50 fleas were then removed from the glass box and placed in a test tube overnight. The following morning all fleas and eggs were removed carefully from the test tube. The feces which had been deposited on the walls of the test tube were taken up in salt solution and injected into 2 guinea pigs. Both of these guinea pigs developed typical clinical endemic typhus. One of these guinea pigs was later found to be immune to a known strain of endemic typhus. The second animal was sacrificed to obtain material for inoculation of other guinea pigs. This strain was carried in animals for four generations, a total of 22 guinea pigs and 2 rabbits being used. Eighteen of these guinea pigs developed typical clinical endemic typhus, and one of these animals, from the fourth transfer generation, was tested for immunity to endemic typhus and found immune. The sera of the two rabbits developed agglutinins for *B. proteus* X₁₉, type O, the serum of one rabbit giving complete agglutination in a dilution of 1:80, while the second showed

complete agglutination at 1:160; incomplete at 1:320 and 1:640; and partial agglutination at 1:1280.

This experiment was repeated twice, the two strains established in these repetitions being known as flea feces virus X-8 and flea feces virus X-13, respectively. Both of these strains were studied carefully in guinea pigs and rabbits for several generations. A total of 51 guinea pigs and 4 rabbits (10 generations) were inoculated with strain flea feces X-8. Thirty-nine of the guinea pigs inoculated with this strain developed clinical endemic typhus, while of the 4 rabbits inoculated, 1 died, and the sera of the 3 remaining developed agglutinins for *B. proteus* X₁₉, type O, as shown in Table 1.

TABLE 1.—Agglutination of *B. proteus* X₁₉, type O, by the sera of rabbits following inoculation with virus strains recovered from feces of typhus-infected fleas

Rabbit	Flea feces X-8									Rabbit	Flea feces X-13								
	Number of weeks after inoculation	Serum dilutions									Number of weeks after inoculation	Serum dilutions							
		10	20	40	80	160	320	640	1,280			10	20	40	80	160	320	640	1,280
4621A-----	0	2	0	0	0	0	0	0	0	4532A-----	0	2	0	0	0	0	0	0	0
	1	4	4	4	3	0	0	0	0		1	4	4	4	4	2	0	0	0
	2	3	4	3	1	0	0	0	0		2	4	4	4	3	2	0	0	0
	3	4	4	4	4	2	0	0	0		3	4	4	4	4	2	0	0	0
											4	4	4	3	0	0	0	0	0
4702A-----	0	0	0	0	0	0	0	0	0	4532B-----	0	3	2	0	0	0	0	0	0
	1	4	4	4	4	2	0	0	0		1	4	4	2	0	0	0	0	0
	2	4	4	4	4	4	3	1	0		2	4	4	4	4	4	3	1	0
	3	4	4	4	4	3	1	0	0		3	4	4	4	4	4	4	3	0
											4	4	4	4	4	4	2	0	0
4702B-----	0	1	0	0	0	0	0	0	0										
	1	3	3	2	1	0	0	0	0										
	2	4	4	4	4	2	0	0	0										
	3	4	4	4	4	2	1	0	0										


Rickettsiae were found readily in smears made from the tunica vaginalis of guinea pigs injected with the flea feces X-8 strain of virus. Of three brains examined histologically, all showed the lesions characteristic of endemic typhus in the guinea pig. That a definite cross immunity existed between this strain of virus and known endemic typhus strains is shown in Charts I and II.

The strain known as flea feces X-13 was studied in guinea pigs and rabbits for nine generations, 66 guinea pigs and 2 rabbits being used. Approximately three-fourths of the guinea pigs developed clinical endemic typhus. The sera of the rabbits developed agglutinins for *B. proteus* X₁₉, type O, as shown in Table 1.

Rickettsiae were found readily in smears made from the tunica vaginalis of guinea pigs infected with this strain of virus. Brains from five guinea pigs from this strain were examined histologically and characteristic lesions of endemic typhus were found in four of them. Clear-cut cross immunity was found to exist between this strain of virus and known strains of endemic typhus virus.

Experimental work on the viability of typhus virus in infected fleas shows that the virus may remain virulent in the rat flea (*Xenopsylla cheopis*) for as long as 36 days after the last infecting feeding. It seems probable that once this species of flea becomes infected it may remain infective through life.

Attempts have been made to recover typhus virus from fleas hatched from eggs of infected fleas. In none of these attempts has




evidence been procured that typhus virus may be transmitted by infected fleas to their offspring through the egg.

In the past few months we have attempted repeatedly to transmit typhus by feeding infected fleas on normal guinea pigs. In these experiments the fleas were confined in test tubes which were closed by stretching chiffon over the mouths of the tubes. The fleas fed readily through the chiffon but in no instance did the guinea pigs develop evidence of typhus, nor were they found later to be immune to subsequent injections of typhus virus.

In view of the negative results in our attempts to transmit typhus by direct bite of infected fleas, arranged in such a manner as to practically eliminate any part the feces might play, we tried to transmit the infection by crushing infected fleas and smearing them on the

abraded abdomen of guinea pigs. In this experiment we were successful.

Without placing too much stress on our negative results in direct feeding of infected fleas, the foregoing work suggests that a probable mechanism by which endemic typhus may be transmitted is through



the rubbing of infected feces into wounds made by the biting of the flea or by scratching.

REFERENCES

- (1) Dyer, R. E., Ceder, E. T., Rumreich, A., and Badger, L. F.: Pub. Health Rep., 46 : 2415 (Oct. 9), 1931. Reprint No. 1517.
- (2) Dyer, R. E., Ceder, E. T., Rumreich, A., and Badger, L. F. : Pub. Health Rep., 46 : 1869 (Aug. 7), 1931. Reprint No. 1498.
- (3) Dyer, R. E., Ceder, E. T., Lillie, R. D., Rumreich, A., and Badger, L. F.: Pub. Health Rep., 46 : 2481 (Oct. 16), 1931. Reprint No. 1520.

ANOPHELES ATROPOS D. & K.—A NEW POTENTIAL CARRIER OF MALARIA ORGANISMS

By BRUCE MAYNE, *Special Expert*, and T. H. D. GRIFFITHS, *Surgeon, United States Public Health Service*

The specimens of the *Anopheles atropos* D. & K. used in the infectivity experiments described here were captured as imagoes on the three days, October 29 and 30 and November 2, 1931, in a salt marsh at Pointe aux Chenes, near Ocean Springs, Miss. It was desirable to supplement these collections with bred-out material, but we were not successful in finding a sufficient number of aquatic forms, due probably to the extreme drought prevailing at this time. Therefore, recourse was had to capturing adults which were attracted to the persons of the collectors. The collections were made by visiting small salt pools deep in the marsh and allowing the mosquitoes to attack while remaining quiet. In this manner two collectors captured approximately 50 female specimens of *Anopheles atropos*, some of which were permitted to become blood engorged. The mosquitoes were collected in glass tubes and transferred immediately to cloth cages, made after the pattern of the Barraud shipping cage. These cages are admirably suited for shipment at long distances, for they are so constructed that the live specimens of mosquitoes are kept in a humid atmosphere by means of moist cotton gauze surrounding the netted fabric protected by the galvanized wire frame.

The specimens while awaiting shipment were maintained by placing partially masticated raisins within reach of the insects. These cages were placed with a final moistening of the gauze pads in stout corrugated cardboard boxes and transported by post to Columbia, S. C. A count of the survivors yielded nearly 100 per cent, showing clearly the advantage of the netted cloth cages of the Barraud type over the metal cloth cages used for comparative purposes.

Table 1 details data in which the specimens of *atropos*, when applied to a suitable carrier of *P. vivax* gametocytes, proved infected on dissection.

TABLE NO. 1.—Designating *atropos* infections

Serial No. of mosquito	Dates of biting carrier	Number of feedings	Date of dissection	Longest possible incubation	Results
	NOVEMBER			Days	
1	1, 5, 9, 11, . . .	4	Nov. 14	14	3 oöcysts, pigmented, largest 16 mu.
2	1, 3, 5, . . .	3	do.	14	19 oöcysts, pigmented, 8-12 mu.
3	1, 3, 5, . . .	3	Nov. 10	10	3 oöcysts, pigmented, maximum 12 mu.
4	1,	1	Nov. 7	7	1 oöcyst, pigmented, 4 by 8 mu.
5	1, 4, 8, 13, . .	4	Nov. 15	15	38 oöcysts, undifferentiated, majority pigmented, 16-35 mu.
6	1, 6, 8, 11, 15, 18.	6	Nov. 19	19	6 oöcysts, 60-64 mu; 2 of them containing sporozoites; others segmented. In addition, 10 oöcysts pigmented, in size from 20-48 mu. No sporozoites in glands.

TABLE NO. 1.—Designating atropos infections—Continued

Serial No. of mosquito	Dates of biting carrier	Number of feedings	Date of dissection	Longest possible incubation	Results
	NOVEMBER—CON.			Days	
7	2, 5, 8, 10...	4	Nov. 14	12	42 oöcysts, all but 1 pigmented, 24-33 mu; average 27 mu; 1 pre-segmented, size 32 mu.
8	2, 5, 8.....	3	Nov. 9	7	60 oöcysts, pigmented, average 8 mu.
9	2, 4.....	2	Nov. 8	6	68 oöcysts, pigmented, average 14 mu; maximum 16 mu.
10	2, 7, 9, 11, 15, 19.	6	Nov. 23	21	2 granulated oöcysts, 20-24 mu; 1 oöcyst capsule. Scanty number of free swimming sporozoites, size 12-13.2 mu. Glands: All lobes swarming with sporozoites; typically active, average size 12 mu, a few at 15.5 mu. Staining characteristic, single and double nucleus. Fields of sporozoites in massed heavy clusters.
12	3.....	1	Nov. 10	7	6 oöcysts, pigmented, 8-14 mu.
13	3, 7, 9.....	3	Nov. 12	9	26 oöcysts, size up to 22 mu; average 16 mu.
14	3, 7, 9.....	3	Nov. 15	12	53 oöcysts, majority pigmented or granulated; size 16-22 mu.
15	4, 6, 8.....	3	Nov. 13	9	24 oöcysts, pigmented; maximum 16 mu, average, 12 mu.
16	4, 6, 8, 10, 15, 21.	6	Nov. 27	23	Gut: More than 3-400 oöcysts covering the blood engorged organ, majority segmented, 12 at least ripe, with sporozoites; many free-moving sporozoites seen. Glands: Packed with very typical sporozoites.
17	4, 7.....	2	Nov. 11	7	2 oöcysts, size 9 mu.
18	4.....	1	Nov. 10	6	3 oöcysts, size 12-16 mu.
19	4, 6, 8, 10...	4	Nov. 14	10	11 oöcysts, pigmented; maximum 23 mu.
20	5, 7, 9, 12, 15, 18, 21.	7	Nov. 25	20	Gut: A few pigmented oöcysts observed, size 17.76 mu. The gut blood engorged. Glands: Negative.
21	5, 9, 11, 15...	4	Nov. 18	13	Approximately 40 oöcysts, 12-28 mu, majority 20-24 mu. Pigmented and granulated.
23	5.....	1	Nov. 8	3	36 oöcysts, size 4-9 mu.
25	6, 9, 12, 15...	4	Nov. 17	11	15 oöcysts, 12-32 mu; pigmented, larger ones granulated.
26	7, 13, 16, 18, 21.	5	Nov. 25	18	Tremendous infection; both stomach and glands containing approximately several hundred oöcysts in various stages of development, particularly mature forms packed with sporozoites; mounting fluid contained matted clusters of actively wriggling sporozoites; thousands of these were observed; thoracic muscles in the region of the glands with extreme numbers of sporozoites; glands heavily packed; size 11-15.5 mu.
27	7.....	1	Nov. 13	6	4 oöcysts, pigmented, 14-16 mu.
29	9, 13.....	2	Nov. 15	6	14 oöcysts, pigmented; average 8 mu.

Summary of Table 1, designating atropos infections

Total dissected.....	28
Total with oöcysts—5 days or more.....	24
Total negative.....	3
Mosquitoes with sporozoites:	
Up to 15 days.....	0
15-23 days—	
Gut with sporozoites.....	4
Gland with sporozoites.....	3
Percentage of infections.....	85.7

SUPPLEMENTARY NOTES TO TABLE NO. 1 ON MOSQUITOES FOUND WITH MATURE ORGANISMS

Specimen No. 6.—This mosquito was induced to bite a patient suffering from the effects of an infection caused by *P. vivax*, resulting from mosquito biting experimentally. Six feedings were obtained during the 19 days' incubation period. The host's blood exhibited on two

occasions as high as 75 mature gametocytes to 1,000 leucocytes counted in a thick smear.

When dissected on November 19 the gut of this mosquito was found heavily engorged with blood undigested from its last meals. There were a total of 16 oöcysts observed, 10 of them 20–48 μ in size, all containing characteristic pigment. Four oöcysts were segmented; pigment here was absent, and the two remaining forms contained sporozoites, probably only recently ripened. The latter oöcysts and the other four just mentioned measured 60–64 μ . A prolonged search failed to produce free sporozoites in the mounting fluid surrounding the gut or in the material from the macerated thorax. The salivary glands appeared quite free of sporozoites.

Specimen No. 10.—Six infective feedings, synchronous with the preceding specimen, were allowed to this mosquito. It survived an incubation period of 21 days. The gut offered as evidence of infection two granulated oöcysts of 20 and 24 microns in size, and one discharged capsule of an oöcyst. Further evidence was observed in the presence of a scanty number of undetached sporozoites. These were 12–13.2 μ and actively motile along the gut wall.

The glands of the dissected mosquito were kept under observation during a period of six hours. All of the six lobes appeared crowded to the maximum capacity with sporozoites, while the forms already liberated in the saline suspension appeared in a swarming mat of typically active organisms. Their movement was undulating, while the tapering ends were observed to curve in the form of a shepherd's crook. The majority were seen with a single nucleus, many with two nuclei. The size varied in length from 12–15.5 μ , the majority measuring 12 μ , and their width being fairly uniform at 1–1.5 μ .

The dissected material was kept at a temperature of 60° F., and there appeared no diminution of activity after six hours.

After staining with Giemsa it was observed that the sporozoites were present in great profusion. They reacted quite specifically to the Giemsa stain. The sporozoites were again measured, the majority appearing contracted in length by 1 micron. They measured 11–14 μ . A single form, apparently unchanged, measured nearly 15.5 μ . It seemed considerably distended and disintegrated.

Specimen No. 16.—This specimen of *atropos* was given an opportunity to become infected during a development of 23 days while it was induced to bite a gametocyte carrier of *P. vivax* on six occasions. This mosquito had been applied to two patients,¹ who were selected for malaria therapy, before it was killed for the purpose of examination.

On the surface of the blood-distended gut wall, on a portion suitable for inspection, there were observed 12 oöcysts of size 55.5 μ ,

¹ Both of these patients showed very marked clinical symptoms of malarial fever with typical specimens of *Plasmodium vivax* in their blood following an incubation period of 13 days and 16 days, respectively.

engorged with sporozoites. Several more oöcysts, 38.4 mu in maximum size, appeared on the edge of the gut tissue in a stage of pre-segmentation. In addition, several oöcyst capsules with collapsed walls were noticed on the gut wall, and after clearing some of the blood from the stomach, it was apparent that the gut surface was fairly covered with oöcysts in a stage of segmentation. There were evidently more than 300 to 400 of these.

Many sporozoites were observed freely moving in the fluid along the gut wall.

The salivary glands appeared packed to the utmost with living sporozoites, showing typical form and behavior when expressed on pressure of the cover glass. They measured in length 11-15.5 mu.

Specimen No. 26.—Five infective feedings were allowed this mosquito. It died after 18 days of parasite development. Upon dissection there was obviously a tremendous invasion of organisms in all stages. The gut contained several hundred oöcysts, particularly of the mature stages. Not only were the oöcysts fairly engorged with live-looking sporozoites, but there were matted clusters of tens of thousands of actively wriggling, sickle-shaped organisms surrounding the alimentary tract in the saline dissecting fluid.

Measurements of some of these oöcysts under usual pressure of cover glass resulted as follows:

Fourteen of the undifferentiated forms appeared to attain a maximum diameter of 66 mu.

Twenty of the segmented forms measured 39.6-50.6 mu.

Twenty of the forms containing sporozoites measured 48.4-61.6 mu.

The undetached sporozoites from the gut wall measured the same size as those examined from the lobes of the salivary glands, namely, 11.10 to 15.54 mu, with an average length of 13.32 mu and a width of 1.4 mu. The glands and the tissue of the macerated thoracic material were unusually heavily infected with great numbers of motile sporozoites measuring as previously recorded.

The controls used for the *atropos* infectivity tests were a collection of anophelines of three species captured from a stable about 20 miles from Columbia. They were treated in the same manner regarding the source of infection and exposure to temperature and humidity as the specimens of *atropos* described in Table 1. These data are described in Table 2.




FIGURE 1.—Top: Portion of wing of *Anopheles walkeri* Theob.; 20, wing of *Anopheles atropos* D. & K.; 22, wing of *Anopheles quadrimaculatus* Say. Reproduced from plates of Howard, Dyar, and Knab. Mosquitoes of North America. Carnegie Press



FIGURE 2.—Photograph of *A. atropos* (X4) specimen No. 16, mentioned in text, showing characteristic *Culex*-like attitude



FIGURE 3.—Portion of gut wall of *atropos* No. 16, showing one ripe oöcyst and ruptured oöcyst capsule




TABLE No. 2.—Designating controls: *Atropos* infections

Species and serial No.	Dates of biting carrier	Number of feedings	Date dissected	Longest possible incubation—days	Results
Quad. M-2....	Oct. 19, 25, 29, and Nov. 2, 5, 8, 11.	7	Nov. 13	25	Several hundreds of oöcysts in all stages up to 65 mu. Sporozoites on gut. Glands: Numerous sporozoites.
Quad. M-4....	Oct. 19, 22, 25, 28.	4	Oct. 31	12	Moderate number of oöcysts; none over 24 mu.
Quad. M-6....	Oct. 19, 23.....	2	Oct. 26	7	14 pigmented oöcysts, size 12 mu and under.
Punct. M-7....	Oct. 19, 22, 25, 30....	4	Nov. 9	20	33 oöcysts up to 65 mu, 6 with sporozoites; many sporozoites free on gut. Glands: maximum number of sporozoites.
Quad. O-2....	Oct. 21, 24.....	2	Oct. 28	7	Several pigmented oöcysts, pigmented up to 20 mu.
Quad. O-3....	Oct. 21, 24, 28.....	3	Oct. 30	9	22 oöcysts, pigmented and presegmented, size 16-24 mu.
Quad. O-7....	Oct. 21, 25, 30, and Nov. 2, 5, 8.	6	Nov. 11	21	More than 100 (majority segmenting) oöcysts, size 45-60 mu. Many free sporozoites seen. Glands: Tremendous sporozoite infection.
Quad. O-9....	Oct. 21, 24, 28, 31, and Nov. 3, 8.	6	Nov. 12	22	Specimen blood engorged at dissection; sporozoites seen along gut wall and in thorax.
Quad. O-10....	Oct. 21, 24, 28, and Nov. 2, 5.	5	Nov. 8	18	Oöcysts; total number 128, 3 with sporozoites; size 45-62-60 mu. Glands: Quite negative.
Punct. O-12....	Oct. 21, 24, 28.....	3	Oct. 30	9	Moderate number of oöcysts, size 16-24 mu, pigmented.
Punct. P-3....	Oct. 22.....	1	Oct. 29	7	11 oöcysts, size 12-20 mu; average 16 mu.
Punct. P-4....	Oct. 22, 25, 28.....	3	Oct. 31	9	Moderate number of oöcysts; maximum size 20 mu.
Quad. P-6....	Oct. 22, 25, 28, and Nov. 1, 4, 8, 11, 13.	8	Nov. 16	24	Approximately 150 oöcysts, 8-68 mu; average about 40 mu. Pigmented, granulated, and segmented forms. Numerous sporozoites in media surrounding stomach. Glands packed with sporozoites. Swarms in fluid active, 12-16 mu in size.
Punct. P-7....	Oct. 22.....	1	Oct. 23	13½	Several hundred ookinetes observed.
Punct. P-9....	do.....	1	Oct. 24	2	A few pigmented forms, quite immature.
Punct. Q-3....	Oct. 23.....	1	Oct. 26	3	Great numbers of pigmented zygotes, less than 8 mu.
Punct. Q-4....	Oct. 23, 26, 30.....	3	Nov. 6	14	About 200 oöcysts pigmented, none reaching segmented stage.
Quad. Q-5....	Oct. 9, 24, and Nov. 3, 6, 11, 15.	6	Nov. 17	23	Upward of 100 oöcysts, 20-72 mu in size; majority 40-48 mu. Numerous sporozoites in mounting fluid. Glands packed with sporozoites, 12-16 mu. in size. Active and typical.
Punct. Q-7....	Oct. 24, 28, 31.....	3	Nov. 3	10	8 oöcysts pigmented, 8-12 mu.
Punct. Q-8....	Oct. 24, 28, 31, and Nov. 6.	4	Nov. 11	18	Approximately 125 oöcysts (100 counted) in all stages, except pigmented, up to 60 mu; majority with sporozoites. Glands: A scanty number of full-sized active sporozoites.
Punct. R-6....	Oct. 30 and Nov. 2, 5, 8, 11.	5	Nov. 12	13	Approximately 125 oöcysts (54 counted); majority 22-28 mu, maximum 32 mu; pre-segmented stage mostly.
Punct. R-9....	Oct. 30.....	1	Nov. 3	4	4 oöcysts, size 8-21 mu.
Punct. R-11....	Oct. 30 and Nov. 2, 7, 10.	4	Nov. 13	14	About 60 oöcysts pigmented, size 5-8 mu.
Punct. R-14....	Nov. 1.....	1	Nov. 8	8	As many as 90 oöcysts pigmented, maximum size 16 mu.
Crucians R-2....	Oct. 26, 30, and Nov. 2, 5.	4	Nov. 10	15	Nearly 200 oöcysts (counted 180); majority stage of segmentation; size 60 mu. Glands: Apparently negative.

Summary of Table 2, designating controls: *Atropos* infections

Total dissected.....	38
Total with oöcysts—5 days or more.....	21
Total negative.....	13
Mosquitoes with sporozoites:	
Up to 15 days.....	0
15 to 25 days—	
Gut with sporozoites.....	1
Gland with sporozoites.....	7
Percentage of infections.....	55.2

There are offered for comparison the results of attempting to infect specimens of *Anopheles quadrimaculatus* obtained in the same general region of the Gulf coast where the specimens of *atropos* were collected.

Six of the specimens which survived the shipment from place of origin and developed the infection after biting the tertian carrier in two to five applications are included in the following table:

TABLE NO. 3.—Regarding infections of *quadrimaculatus*

Serial No. of mosquito	Dates of biting carrier	Number of feedings	Date of dissection	Longest possible incubation	Results
1	Sept. 5, 8, 11, 13.	4	Sept. 28	Days 23	Gut: 9 oöcysts and 5 discharged. Size up to 43 mu granulated and segmenting. Two ripe with sporozoites. Glands: A few sluggishly active sporozoites in mounting fluid. Lobes of glands packed with normal appearing sporozoites, size 12-14 mu.
2	Sept. 5, 8, 11, 13, 15.	5	Oct. 2	27	Gut: Fairly covered with presegmenting oöcysts, size up to 64 mu; average size 45 mu. One with sporozoites. Glands: Devoid of sporozoites.
3	Sept. 5, 8, 11, 13.	4	Sept. 17	12	Gut: 8 oöcysts observed and 2 discharged forms. Glands: Moderate infection; sporozoites quite active and normal.
4	Sept. 5, 8, 12.	3	Sept. 14	9	Gut: 14 oöcysts, 4 of them segmented and sporozoites noted in 2 others. Glands: Scanty number of typical sporozoites. Normal in form, size, and motility. Size 12-14 mu.
5	Sept. 5, 9....	2	Sept. 12	7	Gut: 5 oöcysts present, 24-40 mu, pigmented and granulated forms. Sporozoites absent.
7	Sept. 6, 9, 12.	3	Sept. 15	9	Gut: A total of 152 oöcysts counted, 3 of these 56-68 mu in size. No sporozoites either on gut or in glands.

Summary of Table 3

Total dissected.....	7
Total with oöcysts—5 days or more.....	6
Total negative.....	1
With sporozoites up to 27 days:	
Gut.....	4
Glands.....	3
Percentage of infections.....	85.7

A note on the biological relationships of *Anopheles atropos* is contributed as a supplement to the present experimental data.

Habitat.—In the course of a survey of salt-marsh mosquito-breeding areas of the South Atlantic and Gulf States, conducted by the United States Public Health Service, *Anopheles atropos* has been recorded in the four States of Mississippi, Louisiana, Alabama, and Florida. It is strictly a salt-water mosquito, frequently found in the same habitat as *Aedes sollicitans*, *Ae. taeniorhynchus*, and *Anopheles crucians*.

At Pointe aux Chenes, near Ocean Springs, Miss., where specimens of adults were captured which were employed in the infectivity tests recorded in this paper, are surrounding marshes characteristic of such habitats having a firm alluvial dense root mat formation, covered with a heavy growth of salt grass (*Spartina* spp.). Where salt pools,

which are the favorite production areas of *atropos*, occur in these marshes, the water can scarcely be muddied, the bottom of the pools being sandy, with sides of a firm clay. When production is said to be heavy, larvae of this species inhabit every square foot of water surface.

The preferential breeding place of *A. atropos* is characterized by the junior author as shallow water on muck or alluvial marshes, or in permanent salt pools whose water has a salinity (salinometer with direct salinity reading) of from 3 per cent to 21 per cent.

Host relations.—*Atropos* have been observed in great numbers in occupied rooms in hotels and private homes within flight distance of the production areas. The junior author has personally collected these mosquitoes at such places at Buras, La., and at Biloxi, Miss.

Biting habits.—Close to its breeding place in marsh areas *atropos* is known to attack in direct sunlight as well as by night. It is then a greater torment as a pest than the redoubtable *Aedes sollicitans*, which shares its intrepidity in persisting in its attacks so that one may easily collect it when attached to its host by dislodging it with thumb and finger.

The culexlike attitude of atropos.—*Atropos* is distinguished at once from the common species of anophelines of America by its decided culexlike appearance, especially when attacking or resting after blood engorgement. This is further emphasized by its unorthodox nonanopheline wing, which is clear in the bright sunlight. When observed biting in the direct sunshine, this species assumes the 2-plane angle which does not characterize the common anophelines, namely, *quadrimaculatus*, *punctipennis*, or *crucians*. *Anopheles atropos* is observed to typify less the "standing-on-head" position while biting and often appears "sprawled" when about ready to finish the blood meal. The brown color of the mesonotum, as well as its near *Culex* position, makes this species often mistaken by the unwary for a *Culex*, especially because of its resemblance to *Culex salinarius*.

Morphological characters.—*Anopheles atropos*² is described by the taxonomist as a rather small blackish *Anopheles* with unspotted wings. Its wing scales are entirely dark, not forming spots. Its mesonotum

² The specimens of mosquitoes employed in our experiments were provisionally identified when collected alive and studied while biting and resting. The authors agreed to the specific identification of these specimens as *A. atropos* D. & K. Following the dissection of the stomach and salivary glands, all of the parts that were possible to salvage namely, wings, legs, abdominal integument, thoracic exoskeleton, and head with mouth parts, were meticulously assembled, placed in gelatine capsules, and submitted to Dr. Harold Morrison, Chief of the Taxonomic Division of the U. S. Bureau of Entomology. He, with the assistance of Dr. Alan Stone, dipterist of the U. S. National Museum, courteously consented to attempt to identify the species of the several mosquito remnants submitted. Their report is as follows: Only one of the specimens, namely, No. 14, was found impossible to examine. The remainder were regarded indeterminately, either *Anopheles atropos* Dyar and Knab, or *Anopheles walkeri* Theobald. "*Anopheles atropos* D. & K. can not be distinguished from *A. walkeri* Theob. in the female. It is difficult to distinguish them from *quadrimaculatus* Say. *Atropos* breeds in salt water, *walkeri* in fresh, and both occur in the South. Only a study of the male genitalia will separate these and there is some question as to their specific distinction."

is elongate and deep brown; abdomen blackish in the integument, with dark hairs; legs and palpi entirely dark, the latter with traces of paler markings at the articulations.

Color.—Recently emerged imagoes are very dark, almost a bluish black. Older specimens appear brownish or even remarkably reddish on the mesonotum.

The species of anophelines discussed in this paper can be distinguished in life from its nearest relatives, *Anopheles quadrimaculatus* Say, and *A. walkeri* Theob., but some confusion arises when identification is required of a specimen preserved for the museum. The following parallel, from a description of the females taken from Dyar's Mosquitoes of the Americas (1928), is offered in identifying the two more closely related species, *A. atropos* and *A. walkeri*:

<i>Atropos</i> (female)	<i>Walkeri</i> (female)
Proboscis: straight, black.	Slightly curved, black.
Palpi: black, small faint white rings, bases of joints.	Rather slender, black, yellowish rings at tips of all joints.
Occiput: black, erect forked scales and long bristles, all black.	Black, whitish spot on each side, scales erect, forked, black.
Mesonotum: black, brownish or black hairs; pleurae black.	Dark brown, more or less streaked with whitish; pleurae brown and grayish.
Abdomen: blackish, with brown-black hairs.	Black, with yellowish-brown hairs.
Legs: brown black, without spottings.	Black, with bronzy reflections, femora and tibiae yellowish white at tips.
Wings: scales black, without spots.	Scales black, not or faintly forming spots at bases of second to fourth veins and forks of second and fourth.

Temperature and humidity.—During the 25 days of the experimental investigation the specimens of *Anopheles atropos* and the controls were maintained at a relatively low temperature of 68° to 70° F. during the months of October and November. The relative humidity registered a high mean percentage of 80 to 90.

The conditions maintained for the specimens detailed in Table 3 were a decidedly higher temperature up to the development of sporozoites. The temperature here went to a maximum of 82° F. during the latter part of September and in October, and may account for the great acceleration of the appearance of gland sporozoites, namely, a minimum of nine days. In the other controls of the same species and the same parasite, *P. vivax* sporozoites did not appear before 18 days.

Conclusions.—*Anopheles atropos* D. & K. is presented as a new potential carrier of *Plasmodium vivax*. In infectivity tests it proved equal in efficiency to *Anopheles quadrimaculatus*, *A. crucians*, and *A. punctipennis* used as controls under similar or more favorable conditions.

Acknowledgments

The work of attempted infectivity was conducted at the South Carolina State Hospital, where, through the courtesy of the superintendent, Dr. C. F. Williams, and the medical director, Dr. E. L. Horger, and the other authorities of the State institution, suitable patients were provided for the use of the Government investigators. Mention should be made of the services of Mr. Hans E. Hingst, senior medical technician, who was indefatigable beyond the call of duty in contributing, largely by his skillful dissections, to the success of the experimental procedure.

CURRENT PREVALENCE OF COMMUNICABLE DISEASES IN THE UNITED STATES ¹

November 8-December 5, 1931

The prevalence of certain important communicable diseases, as indicated by weekly telegraphic reports from State health departments to the Public Health Service, is summarized in this report. The underlying statistical data are published weekly in the PUBLIC HEALTH REPORTS under the section entitled "Prevalence of Disease."

Poliomyelitis.—Further recovery from the recent epidemic of poliomyelitis continued through the month of November. For the current 4-week period the number of reported cases was only about 72 per cent of the number reported for the same period last year. The number was, however, more than three times the number of cases recorded for the corresponding period in 1929.

In the New England and Middle Atlantic States, where the epidemic first appeared, the number of cases for the current period was still almost double the number of cases reported for the same period last year. The South Atlantic States compared very favorably with last year and in other regions the decreases in the incidence of the disease ranged from 50 per cent in the West North Central States to 80 per cent in the Mountain and Pacific groups. In the latter group, this period last year marked the first appreciable decrease in the outbreak of poliomyelitis which had begun there earlier in the season. A comparison of this group with 1929, a more nearly normal year, shows that the incidence of the disease during the current period was about 15 per cent in excess of its incidence during the same period in that year.

¹ From the Office of Statistical Investigations, U. S. Public Health Service. The number of States included for the various diseases are as follows: Typhoid fever, 47; poliomyelitis, 48; meningococcus meningitis, 48; smallpox, 48; measles, 45; diphtheria, 47; scarlet fever, 47; influenza, 39 States and New York City. The District of Columbia is counted as a State in these reports.

The total number of cases of poliomyelitis reported for the current 4-week period was 625, approximately 1,200 less than were reported for the preceding 4-week period.

Diphtheria.—The total reported incidence of diphtheria (9,357 cases) for the current period was about 33 per cent higher than that of last year for the same period. All areas contributed to the increase except the New England and Middle Atlantic and East North Central. In the former group a slight decrease (6 per cent) was shown and in the latter group the figure for the current period equaled that of last year. The increases in the various groups ranged from 40 per cent in the Far West groups to 90 per cent in the West North Central group.

For the country as a whole the number of cases reported for the current period was approximately 500 less than was reported for the preceding 4-week period which might indicate that the peak for this year was passed during that period (October 11 to November 7). In each of the two preceding years the peak was reached during the period corresponding to the current 4-week period. For this period in 1930 the reported cases totaled 7,031, and in 1929, 9,405 cases were reported.

Measles.—The usual seasonal increase of measles continued through the current 4-week period. The number of cases (8,805) was about 15 per cent in excess of the number reported for the same period in 1930, but was 10 per cent lower than in 1929. The disease was most prevalent in States along the Atlantic coast, the number of cases being much larger than was reported in either of the two preceding years.

In the New England and Middle Atlantic group the number of cases reported during the current period was 4,993, as compared with 2,900 for the same period last year and 2,711 in 1929. The South Atlantic group reported 980 cases, as compared with 218 in 1930 and 212 in 1929. All other areas showed decreases this year, ranging from 75 per cent in the far west groups to 40 per cent in the Great Lakes region. In 1929 the disease was unusually prevalent in some of these areas, especially the East North Central.

Scarlet fever.—Although the usual seasonal increase in scarlet fever was apparent in all sections of the country, the number of cases (15,281) reported for the current 4-week period came closer to the average for previous years than at any time during the current year. States in the North Central groups showed decreases from last year's figure, but in other areas the increases ranged from 11 per cent to 22 per cent.

Smallpox.—The incidence of smallpox maintained the low level which has prevailed throughout the current year. The reported cases for the current 4-week period numbered 1,124, i. e., about 77 per

cent of the cases recorded for the corresponding period last year and considerably less than one-half of the number in 1929.

Areas showing increases over last year were the New England and Middle Atlantic, West North Central, and South Central. In the New England and Middle Atlantic States the disease continued unusually prevalent in Vermont, and during the week ended December 5 there were 39 cases reported in the State of Connecticut. No cases had been reported from Connecticut since 1929. Out of 449 cases reported during the current period from the West North Central group, Iowa reported 249, as compared with 41 in the same period last year. While the number of cases was not high in the South Central States, it represented a 50 per cent increase over the same period last year.

Meningococcus meningitis.—In relation to previous years the incidence of meningococcus meningitis continued considerably below the level of either of the two preceding years for the period involved. The number of cases reported for the four weeks ended December 5 was 279, as compared with 319 cases for the same period last year and 482 cases in 1929. Each geographic area shared in this favorable decrease except the South Atlantic, where, since almost the beginning of the current year, the incidence has been slightly higher than in either 1930 or 1929.

Typhoid fever.—The incidence of typhoid fever continued to decrease during the 4-week period ended December 5. Compared with previous years the incidence (1,967 cases) was about 12 per cent less than that of last year for the same period but was more than 30 per cent in excess of the incidence in 1929. All areas showed considerable decreases in the numbers of cases occurring during the current period as compared with the preceding 4-week period.

Influenza.—The total number of cases (2,593) reported for the 4-week period ended December 5 was about 65 per cent of the number reported for the same period last year and 50 per cent of the number in 1929. All areas shared in this favorable situation except the West North Central. In that group of States 460 cases were reported as compared with 39 for the same period last year and 65 in 1929. Missouri reported 340 of the 460 cases.

Mortality, all causes.—The mortality from all causes in a group of large cities as summarized by the Bureau of the Census was the lowest in six years, viz., 11.1 per thousand population, annual basis.

COURT DECISION RELATING TO PUBLIC HEALTH

Ordinance relative to closing of barber shops held invalid.—(Mississippi Supreme Court; Knight, Chief of Police, *v.* Johns, 137 So. 509; decided Nov. 2, 1931.) By the terms of an ordinance of the city of

Clarksdale it was made unlawful and punishable by fine and imprisonment "for any barber shop in the said city to open for business before 7.30 in the forenoon and/or to remain open for business after the hour of 6.30 in the afternoon, except that, on week days which immediately precede a holiday, said barber shops may remain open for business until 9 o'clock p. m." The ordinance empowered the city health officer to inspect barber shops, and in one section it was declared that the purpose in prescribing the hours of opening and closing was "to promote the general health and sanitary conditions of the said shops, it being apparent that a better inspection may be had and made between the hours prescribed than at any other time."

The appellee, who owned and operated a barber shop in the city, twice violated the ordinance by keeping his shop open after 6.30 p. m. and was twice arrested. He then secured an injunction restraining the chief of police from further arresting him for violating the ordinance. On appeal, one of the reasons assigned for the validity of the ordinance was that it was designed to fix a reasonable time within which the city inspectors could inspect barber shops in order to ascertain whether the city's sanitary and health ordinances were being obeyed. In holding that the ordinance could not be sustained on this ground, the supreme court said:

The city has the right of inspection reasonably necessary for the enforcement of its health and sanitary ordinances. As we understand the argument, the necessity for the barber-shop-closing ordinance arises because of inconvenience to the city's inspectors of inspecting such shops during the hours the ordinance requires them to be closed. It does not, and could hardly be made to, appear that such inspection must be continuous, covering every hour a barber shop is open; and to compel the closing of barber shops between certain hours, because it will be inconvenient for the city to then inspect them, when they are open at other hours amply sufficient for such inspection, would unnecessarily and unreasonably interfere with the operation thereof.

DEATHS DURING WEEK ENDED DECEMBER 5, 1931

Summary of information received by telegraph from industrial insurance companies for the week ended December 5, 1931, and corresponding week of 1930. (From the Weekly Health Index, issued by the Bureau of the Census, Department of Commerce)

	Week ended Dec. 5, 1931	Corresponding week, 1930
Policies in force.....	74, 178, 223	75, 098, 994
Number of death claims.....	12, 885	13, 993
Death claims per 1,000 policies in force, annual rate.....	9. 1	9. 7
Death claims per 1,000 policies, first 49 weeks of year, annual rate.....	9. 6	9. 5

Deaths¹ from all causes in certain large cities of the United States during the week ended December 5, 1931, infant mortality, annual death rate, and comparison with corresponding week of 1930. (From the Weekly Health Index, issued by the Bureau of the Census, Department of Commerce)

[The rates furnished in this summary are based upon mid-year population estimates derived from the 1930 census.]

City	Week ended Dec. 5, 1931				Corresponding week, 1930		Death rate ² for the first 49 weeks	
	Total deaths	Death rate ¹	Deaths under 1 year	Infant mortality rate ¹	Death rate ¹	Deaths under 1 year	1931	1930
Total (82 cities).....	7,404	10.8	559	44	11.8	731	11.8	11.9
Akron.....	32	6.3	2	20	2.9	3	7.5	7.7
Albany.....	42	17.0	3	60	13.9	1	14.0	14.8
Atlanta.....	65	12.2	2	20	13.4	8	15.0	15.3
White.....	29	8.2	0	0	8.9	5	11.6	11.4
Colored.....	36	20.1	2	58	22.4	3	21.7	23.0
Baltimore.....	189	12.1	19	66	15.7	25	14.2	14.0
White.....	156	12.2	15	67	14.4	15	12.9	12.7
Colored.....	33	11.7	4	64	21.8	10	19.9	19.9
Birmingham.....	68	13.2	4	40	15.5	12	13.1	13.6
White.....	32	10.0	2	34	11.6	3	10.1	10.1
Colored.....	36	18.3	2	49	21.9	9	18.1	19.3
Boston.....	206	13.7	9	26	11.7	26	14.1	14.1
Bridgeport.....	37	13.1	2	34	12.8	5	11.1	10.9
Buffalo.....	117	10.5	12	54	12.9	22	12.8	12.9
Cambridge.....	21	9.6	3	62	11.5	1	12.0	11.8
Camden.....	31	13.6	2	35	11.0	0	14.1	13.4
Canton.....	14	6.8	1	25	9.9	1	10.0	9.9
Chicago.....	590	8.9	48	43	11.4	67	10.5	10.4
Cincinnati.....	128	14.6	7	42	16.0	9	15.7	15.6
Cleveland.....	168	9.6	17	50	10.6	9	11.1	11.1
Columbus.....	68	12.0	3	29	12.9	6	13.5	15.3
Dallas.....	64	12.2	9	10	10.7	8	11.1	11.4
White.....	46	11.1	8	10	9.6	6	9.8	10.5
Colored.....	18	17.6	1	1	16.2	2	17.4	16.2
Dayton.....	47	10.6	6	85	11.8	3	10.5	9.6
Denver.....	82	14.7	9	91	12.5	7	13.8	14.9
Des Moines.....	25	9.0	2	38	10.2	0	10.9	11.6
Detroit.....	222	7.0	24	38	8.5	37	8.1	9.2
Duluth.....	24	12.3	3	81	14.4	2	11.3	11.5
El Paso.....	20	9.9	4	17	17.7	4	15.1	17.0
Erie.....	32	14.2	3	62	5.4	2	10.3	11.0
Fall River.....	27	12.2	1	24	10.0	2	11.1	11.6
Flint.....	15	4.8	2	25	7.6	4	6.8	9.0
Fort Worth.....	36	11.2	3	14	14.3	3	10.5	10.9
White.....	31	11.5	3	14	14.0	3	10.2	10.3
Colored.....	5	9.6	0	1	15.8	0	12.3	13.5
Grand Rapids.....	14	4.3	1	15	10.5	3	9.0	10.1
Houston.....	73	12.3	8	12	13.4	10	11.0	12.1
White.....	45	10.3	4	12	12.0	5	10.2	10.8
Colored.....	28	17.6	4	17	17.3	5	13.5	15.9
Indianapolis.....	88	12.4	5	38	14.3	6	13.6	14.4
White.....	72	11.6	3	26	13.5	6	13.1	13.8
Colored.....	16	18.5	2	123	20.0	0	17.1	21.3
Jersey City.....	59	9.6	8	71	11.0	9	11.2	11.3
Kansas City, Kans.....	28	11.9	1	22	11.5	0	12.6	11.7
White.....	24	12.6	1	27	11.6	0	11.9	11.0
Colored.....	4	8.9	0	0	11.4	0	15.5	14.9
Kansas City, Mo.....	88	11.2	8	64	12.6	4	12.9	13.3
Knoxville.....	31	14.8	4	67	7.8	2	12.5	13.5
White.....	27	15.4	4	98	5.3	2	11.7	12.5
Colored.....	4	11.7	0	0	21.1	0	16.3	18.3
Long Beach.....	25	8.6	0	0	12.0	4	9.8	10.0
Los Angeles.....	263	10.4	7	20	11.4	24	10.6	11.0
Louisville.....	63	10.7	5	45	14.9	12	13.7	13.5
White.....	45	9.0	3	31	13.4	11	12.3	12.1
Colored.....	18	19.7	2	143	23.1	1	21.4	21.6
Lowell.....	31	16.1	1	26	10.4	3	12.8	13.3
Lynn.....	21	10.7	2	58	10.2	3	9.4	10.3
Memphis.....	70	14.1	3	32	15.2	6	16.4	15.9
White.....	34	11.1	1	17	12.3	1	13.4	13.2
Colored.....	36	19.0	2	58	19.9	5	21.3	22.5
Miami.....	20	9.3	0	0	13.2	4	11.6	10.9
White.....	16	9.6	0	0	12.8	4	10.7	9.7
Colored.....	4	8.2	0	0	14.5	0	14.6	15.3

See footnotes at end of table.

Deaths ¹ from all causes in certain large cities of the United States during the week ended December 5, 1931, infant mortality, annual death rate, and comparison with corresponding week of 1930. (From the Weekly Health Index, issued by the Bureau of the Census, Department of Commerce)—Continued

City	Week ended Dec. 5, 1931				Corresponding week, 1930		Death rate ¹ for the first 49 weeks	
	Total deaths	Death rate ²	Deaths under 1 year	Infant mortality rate ³	Death rate ¹	Deaths under 1 year	1931	1930
Milwaukee.....	91	8.0	5	22	12.2	13	9.1	9.6
Minneapolis.....	74	8.1	8	51	11.4	10	10.9	10.7
Nashville ⁴	41	13.7	3	45	14.5	3	16.7	16.5
White.....	21	9.7	1	20	8.9	3	14.3	13.9
Colored.....	20	24.4	2	126	29.2	0	22.9	23.2
New Bedford ⁵	25	11.6	1	26	12.0	3	12.1	11.0
New Haven.....	44	14.1	3	46	6.1	2	12.3	12.6
New Orleans ⁶	122	13.6	15	84	17.2	20	16.6	17.3
White.....	72	11.3	8	68	14.0	14	13.5	14.3
Colored.....	50	19.4	7	116	25.3	6	24.1	24.9
New York.....	1,306	9.6	89	38	10.7	134	11.0	10.7
Bronx Borough.....	185	7.3	9	26	7.3	12	8.1	7.8
Brooklyn Borough.....	445	8.8	32	34	10.0	49	10.1	9.8
Manhattan Borough.....	495	14.2	39	52	15.6	55	16.5	15.9
Queens Borough.....	145	6.6	5	20	8.1	17	7.1	7.0
Richmond Borough.....	36	11.5	4	76	9.2	1	13.5	13.8
Newark, N. J.....	103	12.1	14	74	12.5	8	11.5	12.0
Oakland.....	78	13.9	2	25	11.5	1	10.7	11.0
Oklahoma City.....	41	10.9	5	70	15.8	8	10.6	10.9
Omaha.....	70	16.8	7	81	9.7	6	13.8	13.5
Paterson.....	31	11.6	3	51	13.5	1	13.2	12.1
Peoria.....	20	9.6	1	26	10.9	4	12.4	12.2
Philadelphia.....	477	12.6	26	82	13.7	55	12.9	12.6
Pittsburgh.....	163	12.6	13	45	13.6	12	14.3	13.8
Portland, Oreg.....	74	12.6	1	12	10.5	4	11.6	12.1
Providence.....	53	10.8	4	37	12.6	1	12.6	12.8
Richmond ⁶	41	11.6	7	102	17.6	3	15.3	14.8
White.....	17	6.7	3	66	17.2	2	12.9	12.2
Colored.....	24	23.7	4	173	18.7	1	21.4	21.3
Rochester.....	64	10.1	4	37	8.6	3	11.7	11.5
St. Louis.....	191	12.0	12	43	12.9	10	14.8	14.0
St. Paul.....	64	12.1	7	72	10.0	1	10.4	10.1
Salt Lake City ⁷	31	11.3	2	30	14.1	5	12.0	12.5
San Antonio.....	72	15.6	8	15	15.2	10	14.1	15.8
San Diego.....	57	19.0	0	0	16.7	3	13.6	14.5
San Francisco.....	163	13.1	7	47	12.1	1	12.9	13.0
Schenectady.....	27	14.6	0	0	9.3	1	10.8	11.1
Seattle.....	91	12.8	0	0	10.8	3	11.3	10.8
Somerville.....	17	8.4	1	31	8.0	2	8.7	9.6
South Bend.....	14	6.8	2	52	9.9	2	8.0	9.0
Spokane.....	26	11.7	2	82	14.9	3	12.4	12.8
Springfield, Mass.....	23	7.9	0	0	11.1	2	11.4	12.0
Syracuse.....	41	10.0	5	61	11.9	4	11.5	11.6
Tacoma.....	33	16.0	0	0	7.8	1	12.3	12.4
Toledo.....	71	12.5	8	75	14.5	7	11.8	12.6
Trenton.....	36	15.2	2	37	24.5	4	16.2	16.6
Utica.....	23	11.7	0	0	8.7	2	14.2	14.6
Washington, D. C. ⁸	156	16.6	20	111	15.0	13	15.9	15.3
White.....	85	12.4	5	41	13.4	9	13.6	13.1
Colored.....	71	27.4	15	256	19.2	4	22.1	20.8
Waterbury.....	16	8.3	3	75	12.5	1	9.6	9.5
Wilmington, Del. ⁹	15	7.3	0	0	11.7	2	13.7	14.4
Worcester.....	39	10.3	4	57	10.9	3	12.0	12.6
Yonkers.....	10	3.8	0	0	4.6	1	8.3	8.1
Youngstown.....	26	7.8	2	28	9.2	2	9.9	10.3

¹ Deaths of nonresidents are included. Stillbirths are excluded.

² These rates represent annual rates per 1,000 population, as estimated for 1931 and 1930 by the arithmetical method.

³ Deaths under 1 year of age per 1,000 live births. Cities left blank are not in the registration area for births.

⁴ Data for 77 cities.

⁵ Deaths for week ended Friday.

⁶ For the cities for which deaths are shown by color, the percentages of colored population in 1930 were as follows: Atlanta, 33; Baltimore, 18; Birmingham, 38; Dallas, 17; Fort Worth, 16; Houston, 27; Indianapolis, 12; Kansas City, Kans., 19; Knoxville, 16; Louisville, 15; Memphis, 38; Miami, 23; Nashville, 28; New Orleans, 29; Richmond, 29; and Washington, D. C., 27.

⁷ Population Apr. 1, 1930; decreased 1920 to 1930, no estimate made.

PREVALENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

CURRENT WEEKLY STATE REPORTS

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers

Reports for Weeks Ended December 12, 1931, and December 13, 1930

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended December 12, 1931, and December 13, 1930

Division and State	Diphtheria		Influenza		Measles		Meningococcus meningitis	
	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930
New England States:								
Maine.....	7	4		1	264	24	0	0
New Hampshire.....	7	2			5		0	0
Vermont.....	1	5			87	11	0	0
Massachusetts.....	66	93	5	9	180	232	2	2
Rhode Island.....	7	16	3		338	2	0	0
Connecticut.....	5	17	3	1	53	105	1	3
Middle Atlantic States:								
New York.....	124	97	111	113	401	209	8	17
New Jersey.....	44	70	11	16	34	118	6	2
Pennsylvania.....	120	138			625	381	10	3
East North Central States:								
Ohio.....	118	98	22	25	124	57	2	5
Indiana.....	72	71	22	2	30	119	6	4
Illinois.....	161	179	73	29	34	253	5	11
Michigan.....	52	81	11	1	87	89	4	7
Wisconsin.....	23	17	19	21	57	206	1	3
West North Central States:								
Minnesota.....	26	15	1		11	11	0	1
Iowa.....	21	7			2	5	1	0
Missouri.....	90	53	7	9	5	554	1	10
North Dakota.....	30	5			16	5	0	0
South Dakota.....	8	5			125	2	0	2
Nebraska.....	17	15			22	1	0	2
Kansas.....	75	34			24	2	1	2
South Atlantic States:								
Delaware.....	14	3		2	2		0	0
Maryland ¹	70	40	16	22	6	8	0	1
District of Columbia ¹	15	19	2		2	3	1	0
Virginia.....							1	
West Virginia.....	53	27	5	32	286	12	1	2
North Carolina ¹	87	89	32	26	19	44	3	3
South Carolina.....	13	29	406	625	13		0	4
Georgia ¹	32	52	67	88	2	37	1	1
Florida ¹	16	15	2		2	12	1	0

¹ New York City only.

² Week ended Friday.

³ Typhus fever, 1931, 8 cases: 1 case in District of Columbia, 1 case in North Carolina, 2 cases in Georgia, 3 cases in Florida, and 1 case in Alabama.

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended December 12, 1931, and December 13, 1930—Continued

Division and State	Diphtheria		Influenza		Measles		Meningococcus meningitis	
	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930
East South Central States:								
Kentucky.....	94	17					4	1
Tennessee.....	66	29	37	60	8	51	3	3
Alabama ¹	84	82	21	52	18	148	3	6
Mississippi.....	51	29					0	1
West South Central States:								
Arkansas.....	30	12	11	29	13	2	1	0
Louisiana.....	37	38	27	5		3	0	5
Oklahoma ²	97	59	47	45	1	30	0	0
Texas.....	266	56	14	53	3	54	2	0
Mountain States:								
Montana.....	1	2	1		177		0	0
Idaho.....	1					5	0	2
Wyoming.....	7	1		6	1		0	1
Colorado.....	2	11			3	49	1	3
New Mexico.....	14	9			3	38	0	0
Arizona.....	14	4	7	5	4	59	2	3
Utah ³	2	2	3	8	4	1	1	2
Pacific States:								
Washington.....	5	12			57	22	1	2
Oregon.....	1	10	18	17	12	29	0	2
California.....	81	56	105	50	146	221	8	5

Division and State	Poliomyelitis		Scarlet fever		Smallpox		Typhoid fever	
	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930
New England States:								
Maine.....	0	2	44	15	0	0	3	4
New Hampshire.....	0	0	15	2	0	0	0	0
Vermont.....	2	0	8	7	7	0	0	1
Massachusetts.....	7	6	300	236	0	0	3	9
Rhode Island.....	1	0	18	33	0	0	0	0
Connecticut.....	4	0	48	59	15	0	1	5
Middle Atlantic States:								
New York.....	11	4	432	511	40	9	25	26
New Jersey.....	3	0	111	182	0	0	4	2
Pennsylvania.....	7	1	414	451	1	0	26	34
East North Central States:								
Ohio.....	2	11	516	547	13	53	19	23
Indiana.....	1	1	143	189	8	71	12	4
Illinois.....	13	5	367	388	19	36	19	27
Michigan.....	3	3	188	228	14	34	5	13
Wisconsin.....	5	2	89	121	10	18	1	3
West North Central States:								
Minnesota.....	8	2	40	71	6	11	1	1
Iowa.....	3	4	47	53	41	14	1	1
Missouri.....	2	0	74	93	6	5	4	4
North Dakota.....	0	0	22	25	2	5	1	1
South Dakota.....	0	4	16	11	10	12	1	1
Nebraska.....	0	3	27	38	6	7	2	1
Kansas.....	1	3	68	51	5	25	3	2
South Atlantic States:								
Delaware.....	0	0	7	22	0	0	1	0
Maryland ⁴	1	0	109	76	0	0	6	9
District of Columbia ⁴	0	0	21	29	0	0	1	0
Virginia.....								
West Virginia.....	0	0	46	57	4	23	21	15
North Carolina ⁴	0	1	85	82	0	1	6	3
South Carolina.....	0	0	15	20	0	0	9	24
Georgia ⁴	1	0	35	49	2	0	14	9
Florida ⁴	0	0	9	5	2	1	3	0

¹ Week ended Friday.

² Typhus fever, 1931, 8 cases: 1 case in District of Columbia, 1 case in North Carolina, 2 cases in Georgia, 3 cases in Florida, and 1 case in Alabama.

⁴ Figures for 1931 are exclusive of Oklahoma City and Tulsa.

Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended December 12, 1931, and December 13, 1930—Continued

Division and State	Polliomyelitis		Scarlet fever		Smallpox		Typhoid fever	
	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930	Week ended Dec. 12, 1931	Week ended Dec. 13, 1930
East South Central States:								
Kentucky.....	2	0	78	25	0	8	16	1
Tennessee.....	0	1	53	51	3	2	14	3
Alabama ¹	8	0	60	86	0	0	28	22
Mississippi.....	0	0	24	33	4	1	6	10
West South Central States:								
Arkansas.....	0	2	23	17	7	0	14	16
Louisiana.....	0	0	22	24	3	14	33	20
Oklahoma ¹	3	2	38	34	2	21	11	9
Texas.....	0	3	71	47	7	16	20	6
Mountain States:								
Montana.....	3	0	47	42	1	14	0	2
Idaho.....	0	0	5	1	0	1	0	0
Wyoming.....	0	0	11	21	0	0	0	1
Colorado.....	0	2	40	62	0	4	2	1
New Mexico.....	0	1	9	11	0	2	9	16
Arizona.....	0	0	5	5	0	0	0	4
Utah ¹	0	0	12	6	0	0	0	0
Pacific States:								
Washington.....	3	1	66	45	15	25	7	5
Oregon.....	0	1	18	22	6	19	6	4
California.....	3	15	163	99	5	46	10	4

¹ Week ended Friday.

² Typhus fever, 1931, 8 cases: 1 case in District of Columbia, 1 case in North Carolina, 2 cases in Georgia, 3 cases in Florida, and 1 case in Alabama.

³ Figures for 1931 are exclusive of Oklahoma City and Tulsa.

SUMMARY OF MONTHLY REPORTS FROM STATES

The following summary of cases reported monthly by States is published weekly and covers only those States from which reports are received during the current week:

State	Menin- gococ- cus menin- gitis	Diph- theria	Influ- enza	Ma- laria	Measles	Pel- lagra	Polio- mye- litis	Scarlet fever	Small- pox	Ty- phoid fever
<i>October, 1931</i>										
Kansas.....	5	217	4	2	59		2	275	11	45
<i>November, 1931</i>										
District of Columbia.....	2	60	2		9		0	92	0	14
Iowa.....	7	83			13		37	201	258	16
Maine.....	2	17	4		782		13	139	0	16
Massachusetts.....	12	243	19	4	390	5	56	906	0	15
Nebraska.....		93	16		42		4	108	29	5
New Hampshire.....		21					2	23		1
Vermont.....		30			141		8	58	75	0
Wyoming.....					6		0	31	2	1

October, 1931		Lethargic encephalitis:	
Kansas:	Cases	Massachusetts.....	Cases
Actinomycosis.....	1	Mumps:	3
Chicken pox.....	175	Iowa.....	14
German measles.....	4	Maine.....	10
Impetigo contagiosa.....	22	Massachusetts.....	627
Mumps.....	88	Nebraska.....	43
Ptomaine poisoning.....	1	Vermont.....	53
Scabies.....	23	Wyoming.....	20
Septic sore throat.....	4	Ophthalmia neonatorum:	
Tetanus.....	1	Massachusetts.....	96
Trench mouth.....	1	Rabies in animals:	
Tularæmia.....	1	Vermont.....	1
Undulant fever.....	2	Septic sore throat:	
Vincent's angina.....	10	Iowa.....	1
Whooping cough.....	62	Maine.....	2
		Massachusetts.....	21
November, 1931		Tetanus:	
Anthrax:		Maine.....	1
Massachusetts.....	1	Trachoma:	
Nebraska.....	1	Massachusetts.....	5
Chicken pox:		Trichinosis:	
District of Columbia.....	22	Massachusetts.....	2
Iowa.....	363	Undulant fever:	
Maine.....	193	Iowa.....	7
Massachusetts.....	488	Massachusetts.....	8
Nebraska.....	165	Vermont.....	1
Vermont.....	253	Vincent's angina:	
Wyoming.....	31	Iowa.....	8
Conjunctivitis:		Maine.....	4
Maine.....	2	Whooping cough:	
Dysentery:		District of Columbia.....	67
Iowa.....	1	Iowa.....	111
Massachusetts.....	5	Maine.....	80
German measles:		Massachusetts.....	474
Iowa.....	6	Nebraska.....	52
Massachusetts.....	66	Vermont.....	277
Impetigo contagiosa:		Wyoming.....	18
Iowa.....	3		
Lead poisoning:			
Massachusetts.....	6		

ADMISSIONS TO HOSPITALS FOR THE INSANE, AUGUST, 1929

Reports for the month of August, 1929, showing new admissions to hospitals for the care and treatment of the insane, were received by the Public Health Service from 115 hospitals, located in 39 States, the District of Columbia, and the Territory of Hawaii. The 115 hospitals had 180,155 patients on August 31, 1929—95,488 males and 84,667 females, 113 males per 100 females.

The following table shows the number of new admissions for the month of August, 1929, by psychoses:

Psychoses	Number of first admissions		
	Male	Female	Total
1. Traumatic psychoses.....	17	2	19
2. Senile psychoses.....	165	145	310
3. Psychoses with cerebral arteriosclerosis.....	186	127	313
4. General paralysis.....	236	75	311
5. Psychoses with cerebral syphilis.....	22	10	32
6. Psychoses with Huntington's chorea.....	1	2	3
7. Psychoses with brain tumor.....	1	1	2
8. Psychoses with other brain or nervous disease.....	27	11	38
9. Alcoholic psychoses.....	125	12	137
10. Psychoses due to drugs and other exogenous toxins.....	16	9	25
11. Psychoses with pellagra.....	12	24	36
12. Psychoses with other somatic diseases.....	43	55	98
13. Manic-depressive psychoses.....	174	243	417
14. Involution melancholia.....	18	48	66
15. Dementia præcox (schizophrenia).....	310	286	596
16. Paranoia and paranoid conditions.....	28	30	58
17. Epileptic psychoses.....	42	29	71
18. Psychoneuroses and neuroses.....	22	52	74
19. Psychoses with psychopathic personality.....	23	9	32
20. Psychoses with mental deficiency.....	63	63	126
21. Undiagnosed psychoses.....	164	103	267
22. Without psychosis.....	184	46	230
Total.....	1,879	1,382	3,261

During the month of August, 1929, there were 3,261 new admissions to the hospitals, 57.6 per cent of these being males and 42.4 per cent females—136 males per 100 females. Four hundred and ninety-seven of the new admissions were reported as undiagnosed or "without psychosis." There were 2,764 new admissions for whom a provisional diagnosis was made. Of these 2,764 patients, cases of dementia præcox constituted 21.6 per cent; manic-depressive psychoses, 15.1 per cent; psychoses with cerebral arteriosclerosis, 11.3 per cent; general paralysis, 11.3 per cent; and senile psychoses, 11.2 per cent. These five classes accounted for 70.4 per cent of the new admissions for which a diagnosis was given.

The following table shows the number of patients in the hospitals and on parole on August 31, 1929:

	Total patients on books		
	Male	Female	Total
Total patients on books last day of month:			
In hospitals.....	85,443	76,644	162,087
On parole or otherwise absent, but still on books.....	10,045	8,023	18,068
Total.....	95,488	84,667	180,155

Of the 180,155 patients, 10,045 males and 8,023 females were on parole or otherwise absent but still on the books at the end of the month—10.5 per cent of the males, 9.5 per cent of the females, and 10.0 per cent of the total number of patients.

GENERAL CURRENT SUMMARY AND WEEKLY REPORTS FROM CITIES

The 96 cities reporting cases used in the following table are situated in all parts of the country and have an estimated aggregate population of more than 33,360,000. The estimated population of the 89 cities reporting deaths is more than 31,815,000. The estimated expectancy is based on the experience of the last nine years, excluding epidemics.

Weeks ended December 5, 1931, and December 6, 1930

	1931	1930	Estimated expectancy
<i>Cases reported</i>			
Diphtheria:			
46 States.....	2,288	1,666	
96 cities.....	643	560	1,016
Measles:			
45 States.....	2,796	2,896	
96 cities.....	721	894	
Meningococcus meningitis:			
46 States.....	81	105	
96 cities.....	41	37	
Poliomyelitis:			
46 States.....	94	108	
Scarlet fever:			
46 States.....	3,766	3,889	
96 cities.....	1,145	1,270	1,083
Smallpox:			
46 States.....	316	616	
96 cities.....	33	44	23
Typhoid fever:			
46 States.....	416	407	
96 cities.....	47	63	41
<i>Deaths reported</i>			
Influenza and pneumonia:			
89 cities.....	585	650	
Smallpox:			
89 cities.....	0	0	

City reports for week ended December 5, 1931

The "estimated expectancy" given for diphtheria, poliomyelitis, scarlet fever, smallpox, and typhoid fever is the result of an attempt to ascertain from previous occurrence the number of cases of the disease under consideration that may be expected to occur during a certain week in the absence of epidemics. It is based on reports to the Public Health Service during the past nine years. It is in most instances the median number of cases reported in the corresponding weeks of the preceding years. When the reports include several epidemics, or when for other reasons the median is unsatisfactory, the epidemic periods are excluded, and the estimated expectancy is the mean number of cases reported for the week during nonepidemic years.

If the reports have not been received for the full nine years, data are used for as many years as possible, but no year earlier than 1922 is included. In obtaining the estimated expectancy, the figures are smoothed when necessary to avoid abrupt deviation from the usual trend. For some of the diseases given in the table the available data were not sufficient to make it practicable to compute the estimated expectancy.

Division, State, and city	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
		Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
NEW ENGLAND								
Maine:								
Portland.....	8	1	2	-----	0	21	0	1
New Hampshire:								
Concord.....	0	0	0	-----	0	0	0	1
Nashua.....	0	0	0	-----	0	0	1	0
Vermont:								
Barre.....	0	0	0	-----	0	1	0	0
Burlington.....	3	1	0	-----	0	10	0	0
Massachusetts:								
Boston.....	64	38	17	1	0	3	10	17
Fall River.....	9	4	2	-----	0	1	1	3
Springfield.....	7	5	0	-----	0	4	8	1
Worcester.....	8	6	0	-----	0	1	61	1
Rhode Island:								
Pawtucket.....	0	2	0	-----	0	0	0	0
Providence.....	4	9	1	2	0	169	12	2
Connecticut:								
Bridgeport.....	2	5	0	3	1	0	0	5
Hartford.....	4	6	2	-----	0	0	16	2
New Haven.....	16	1	0	4	0	0	7	5
MIDDLE ATLANTIC								
New York:								
Buffalo.....	47	15	6	1	1	2	0	12
New York.....	138	173	92	21	3	43	25	120
Rochester.....	17	4	0	-----	0	20	8	4
Syracuse.....	11	2	0	-----	0	3	4	2
New Jersey:								
Camden.....	3	7	5	2	0	0	0	7
Newark.....	22	16	2	3	0	0	8	7
Trenton.....	6	2	2	-----	0	0	4	1
Pennsylvania:								
Philadelphia.....	118	60	6	6	3	2	20	40
Pittsburgh.....	64	22	7	1	3	177	52	18
Reading.....	20	2	0	-----	0	1	0	2
EAST NORTH CENTRAL								
Ohio:								
Cincinnati.....	16	12	12	-----	1	0	0	12
Cleveland.....	171	38	4	6	1	23	82	17
Columbus.....	22	7	8	-----	0	3	6	1
Toledo.....	79	8	6	-----	0	1	0	5
Indiana:								
Fort Wayne.....	2	5	11	-----	0	0	0	0
Indianapolis.....	60	12	6	-----	2	4	53	10
South Bend.....	2	2	0	-----	0	0	0	0
Terre Haute.....	9	1	2	-----	0	0	0	2
Illinois:								
Chicago.....	106	121	74	5	5	15	9	33
Peoria.....	14	2	6	-----	0	0	1	1
Springfield.....	2	2	3	-----	0	0	1	1
Michigan:								
Detroit.....	44	60	30	2	1	1	2	17
Flint.....	24	3	1	-----	0	3	12	0
Grand Rapids.....	21	1	0	-----	0	1	5	0

City reports for week ended December 5, 1931—Continued

Division, State, and city	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases reported	Mumps, cases reported	Pneumonia, deaths reported
		Cases, estimated expectancy	Cases reported	Cases reported	Deaths reported			
EAST NORTH CENTRAL—Con.								
Wisconsin:								
Kenosha.....	4	1	0	-----	0	0	2	0
Madison.....	8	1	7	-----	-----	1	1	-----
Milwaukee.....	77	15	4	-----	0	1	16	0
Racine.....	41	2	0	-----	0	0	26	0
Superior.....	2	1	0	-----	0	0	8	0
WEST NORTH CENTRAL								
Minnesota:								
Duluth.....	19	0	0	-----	0	0	0	1
Minneapolis.....	74	20	14	-----	1	1	39	4
St. Paul.....	17	7	1	1	1	0	4	3
Iowa:								
Davenport.....	3	1	1	-----	-----	0	0	-----
Des Moines.....	1	2	6	-----	-----	0	0	-----
Sioux City.....	18	1	4	-----	-----	1	1	-----
Waterloo.....	12	0	1	-----	-----	0	0	-----
Missouri:								
Kansas City.....	30	8	9	-----	0	1	0	6
St. Joseph.....	6	2	6	-----	0	1	1	2
St. Louis.....	23	43	38	-----	-----	3	2	3
North Dakota:								
Fargo.....	23	0	0	-----	0	0	0	1
Grand Forks.....	4	0	0	-----	-----	0	0	-----
South Dakota:								
Aberdeen.....	20	0	0	-----	-----	37	0	-----
Sioux Falls.....	0	0	0	-----	-----	0	0	-----
Nebraska:								
Omaha.....	30	9	31	-----	0	2	5	5
Kansas:								
Topeka.....	7	1	1	-----	0	1	0	3
Wichita.....	15	2	11	-----	0	4	0	2
SOUTH ATLANTIC								
Delaware:								
Wilmington.....	1	2	0	-----	0	0	1	0
Maryland:								
Baltimore.....	71	24	10	4	0	4	26	18
Cumberland.....	12	0	0	-----	0	1	0	0
Frederick.....	0	0	0	-----	0	0	0	1
District of Columbia:								
Washington.....	4	18	20	1	1	2	0	16
Virginia:								
Lynchburg.....	-----	4	-----	-----	-----	-----	-----	-----
Norfolk.....	5	3	4	-----	0	0	0	0
Richmond.....	4	14	21	-----	0	0	0	4
Roanoke.....	8	3	6	-----	0	0	0	0
West Virginia:								
Charleston.....	7	2	6	-----	0	0	0	3
Huntington.....	0	-----	2	-----	0	0	0	0
Wheeling.....	3	1	0	-----	0	0	0	2
North Carolina:								
Raleigh.....	1	2	4	-----	0	15	0	1
Wilmington.....	0	2	0	-----	0	0	0	2
Winston-Salem.....	13	3	2	-----	0	0	0	2
South Carolina:								
Charleston.....	1	1	2	25	0	0	0	2
Columbia.....	0	1	1	-----	0	0	0	14
Greenville.....	1	0	0	-----	0	0	0	0
Georgia:								
Atlanta.....	18	7	5	5	0	0	1	7
Brunswick.....	0	0	0	-----	0	0	3	0
Savannah.....	0	2	1	31	1	0	0	1
Florida:								
Miami.....	0	2	2	-----	0	0	0	1
Tampa.....	0	2	1	-----	0	0	0	0

City reports for week ended December 5, 1931—Continued

Division, State, and city	Chicken pox, cases reported	Diphtheria		Influenza		Measles, cases re- ported	Mumps, cases re- ported	Pneu- monia, deaths reported
		Cases, estimated expect- ancy	Cases reported	Cases reported	Deaths reported			
EAST SOUTH CENTRAL								
Kentucky:								
Covington.....	0	1	0		0	0	0	0
Lexington.....	1		1		0	0	1	0
Louisville.....	6		2		0	0	0	4
Tennessee:								
Memphis.....	2	8	15		1	1	1	4
Nashville.....	0	3	4		3	0	0	2
Alabama:								
Birmingham.....	1	7	8		1	2	0	9
Mobile.....	0	1	0		1	0	0	0
Montgomery.....	1	2	1	4		3	4	
WEST SOUTH CENTRAL								
Arkansas:								
Fort Smith.....	0	1	3			0	0	
Little Rock.....	1	1	7		0	0	1	2
Louisiana:								
New Orleans.....	0	15	10	1	0	0	0	12
Shreveport.....	7	1	2		0	7	0	4
Oklahoma:								
Muskogee.....	1		5		0	0	0	0
Texas:								
Dallas.....	2	18	17	1	1	0	0	9
Fort Worth.....	1	11	21		0	0	0	5
Galveston.....	0	1	5		0	0	0	2
Houston.....	0	10	28		0	1	0	7
San Antonio.....	0	5	0		1	0	0	3
MOUNTAIN								
Montana:								
Billings.....	0	0	0		0	60	0	0
Great Falls.....	0	0	0		0	1	0	0
Helena.....	1	0	0		0	14	0	0
Missoula.....	0	0	0		0	0	0	0
Idaho:								
Boise.....	0	0	0		0	0	0	1
Colorado:								
Denver.....	35	10	6		1	3	0	8
Pueblo.....	16	1	0		0	0	0	1
New Mexico:								
Albuquerque.....	7	1	0		0	1	1	1
Arizona:								
Phoenix.....	0	0	1		0	1	0	1
Utah:								
Salt Lake City.....	90	4	0		0	0	3	3
Nevada:								
Reno.....	0	0	0		0	0	0	1
PACIFIC								
Washington:								
Seattle.....	79	5	5			32	22	
Spokane.....	8	2	0			1	0	
Tacoma.....	19	3	3		0	0	2	3
Oregon:								
Portland.....	28	11	0	4	0	5	16	3
Salem.....	6	1	0	7	0	0	2	0
California:								
Los Angeles.....	31	38	33	41	8	8	11	11
Sacramento.....	3	3	1	1	1	44	0	8
San Francisco.....	62	14	3	9	2	7	3	10

City reports for week ended December 5, 1931—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culo- sis, deaths reported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
NEW ENGLAND											
Maine:											
Portland.....	3	12	0	0	0	1	0	0	0	4	27
New Hampshire:											
Concord.....	0	1	0	0	0	0	0	0	0	0	18
Nashua.....	0	0	0	0	0	0	0	0	0	1	-----
Vermont:											
Barre.....	0	0	0	0	0	2	0	0	0	3	5
Burlington.....	1	0	0	0	0	1	0	0	0	0	7
Massachusetts:											
Boston.....	64	60	0	0	0	14	2	1	0	28	206
Fall River.....	3	5	0	0	0	1	0	0	0	1	27
Springfield.....	5	2	0	0	0	0	0	0	1	6	29
Worcester.....	12	26	0	0	0	3	0	0	0	11	39
Rhode Island:											
Pawtucket.....	1	0	0	0	0	0	0	0	0	0	-----
Providence.....	11	9	0	0	0	4	0	0	0	3	53
Connecticut:											
Bridgeport.....	6	4	0	23	0	2	0	1	0	0	37
Hartford.....	6	2	0	0	0	0	0	0	0	4	41
New Haven.....	3	1	0	0	0	0	0	0	0	8	44
MIDDLE ATLANTIC											
New York:											
Buffalo.....	22	24	0	1	0	8	1	1	0	36	115
New York.....	124	114	0	0	0	83	14	6	2	93	1,306
Rochester.....	8	37	0	0	0	2	0	0	0	9	57
Syracuse.....	9	23	0	0	0	0	0	0	0	57	41
New Jersey:											
Camden.....	4	5	0	0	0	1	0	0	0	0	31
Newark.....	13	17	0	0	0	8	0	0	0	32	103
Trenton.....	3	3	0	0	0	7	0	1 ²	0	1	36
Pennsylvania:											
Philadelphia.....	69	71	0	0	0	31	3	1	0	120	477
Pittsburgh.....	39	53	0	1	0	5	0	2	1	25	163
Reading.....	0	0	0	0	0	1	0	0	0	2	29
EAST NORTH CENTRAL											
Ohio:											
Cincinnati.....	17	49	0	0	0	11	1	0	0	2	128
Cleveland.....	34	54	0	0	0	21	1	0	0	126	168
Columbus.....	10	20	0	0	0	2	0	0	0	5	68
Toledo.....	12	8	1	0	0	1	1	5	0	47	71
Indiana:											
Fort Wayne.....	3	1	0	0	0	1	0	0	0	0	27
Indianapolis.....	14	8	3	0	0	5	0	0	0	9	-----
South Bend.....	3	2	0	0	0	0	0	0	0	1	14
Terre Haute.....	3	1	0	0	0	0	0	0	0	0	19
Illinois:											
Chicago.....	100	125	0	0	0	37	3	3	0	144	580
Peoria.....	-----	5	-----	0	0	1	-----	0	0	11	20
Springfield.....	2	11	0	0	0	0	0	0	0	8	27
Michigan:											
Detroit.....	82	73	0	0	0	20	1	3	0	46	222
Flint.....	11	4	1	0	0	2	0	0	0	20	15
Grand Rapids.....	10	6	1	0	0	0	0	0	0	2	14
Wisconsin:											
Kenosha.....	0	4	1	0	0	0	0	0	0	5	6
Madison.....	2	1	1	0	-----	-----	0	0	-----	0	-----
Milwaukee.....	19	16	0	0	0	0	1	0	0	93	91
Racine.....	5	3	0	0	0	2	0	0	0	0	13
Superior.....	3	0	0	0	0	0	0	0	0	0	5

¹ Nonresidents.

City reports for week ended December 5, 1931—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culo- sis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
WEST NORTH CENTRAL											
Minnesota:											
Duluth.....	9	1	0	0	0	0	0	1	0	0	24
Minneapolis.....	41	14	1	0	0	2	0	0	0	15	74
St. Paul.....	17	16	0	0	0	1	0	0	0	3	73
Iowa:											
Davenport.....	1	2	2	0	0	0	0	0	0	0	0
Des Moines.....	9	10	2	0	0	0	0	0	0	0	25
Sioux City.....	2	1	0	1	0	0	0	0	0	7	0
Waterloo.....	3	1	0	0	0	0	0	1	0	11	0
Missouri:											
Kansas City.....	14	19	0	0	0	4	0	0	0	6	88
St. Joseph.....	3	1	0	0	0	0	0	0	1	3	29
St. Louis.....	36	15	0	0	0	11	2	0	0	51	191
North Dakota:											
Fargo.....	2	4	0	0	0	0	0	0	0	4	10
Grand Forks.....	1	1	0	0	0	0	0	0	0	0	0
South Dakota:											
Aberdeen.....	0	2	0	0	0	0	0	0	0	5	0
Sioux Falls.....	1	0	0	0	0	0	0	0	0	0	7
Nebraska:											
Omaha.....	7	9	2	0	0	2	0	0	0	1	70
Kansas:											
Topeka.....	2	3	1	0	0	0	0	0	0	6	13
Wichita.....	4	0	0	1	0	1	0	0	0	0	31
SOUTH ATLANTIC											
Delaware:											
Wilmington.....	2	0	0	0	0	0	0	0	0	4	15
Maryland:											
Baltimore.....	22	18	0	0	0	10	2	3	0	101	189
Cumberland.....	1	4	0	0	0	0	0	0	0	2	12
Frederick.....	0	2	0	0	0	0	0	0	0	0	4
District of Colum- bia:											
Washington.....	18	16	0	0	0	9	1	0	0	14	166
Virginia:											
Lynchburg.....	1	0	0	0	0	0	0	0	0	0	0
Norfolk.....	3	11	0	0	0	1	0	0	0	0	0
Richmond.....	8	20	0	0	0	3	0	0	0	2	49
Roanoke.....	4	2	0	0	0	0	0	0	1	1	13
West Virginia:											
Charleston.....	2	1	0	0	0	0	0	12	1	3	22
Huntington.....	2	6	0	0	0	0	0	0	0	0	0
Wheeling.....	2	1	0	0	0	1	0	0	0	4	20
North Carolina:											
Raleigh.....	3	1	0	0	0	0	0	0	0	3	11
Wilmington.....	1	0	0	0	0	1	0	0	0	4	13
Winston-Salem.....	2	2	1	0	0	1	0	0	0	7	15
South Carolina:											
Charleston.....	2	1	0	0	0	3	0	0	0	0	20
Columbia.....	1	0	0	0	0	1	0	0	1	0	61
Greenville.....	0	1	0	0	0	0	0	0	0	0	0
Georgia:											
Atlanta.....	6	12	0	0	0	6	0	0	0	0	65
Brunswick.....	0	0	0	0	0	1	0	0	0	0	5
Savannah.....	1	2	0	0	0	0	1	0	0	0	30
Florida:											
Miami.....	1	0	0	0	0	2	0	0	0	0	20
Tampa.....	1	5	0	0	0	0	0	3	0	0	19
EAST SOUTH CENTRAL											
Kentucky:											
Covington.....	4	2	0	0	0	0	0	0	0	0	12
Lexington.....	2	2	0	0	0	0	0	0	0	1	15
Louisville.....	16	0	0	0	0	2	0	0	0	14	63
Tennessee:											
Memphis.....	7	8	1	0	0	6	2	1	1	20	70
Nashville.....	4	3	0	0	0	3	1	0	0	7	41
Alabama:											
Birmingham.....	4	6	0	0	0	3	1	1	0	1	68
Mobile.....	1	3	0	0	0	4	0	0	0	0	31
Montgomery.....	0	0	0	0	0	0	0	0	0	0	0

* Nonresident.

City reports for week ended December 5, 1931—Continued

Division, State, and city	Scarlet fever		Smallpox			Tuber- culo- sis, deaths re- ported	Typhoid fever			Whoop- ing cough, cases re- ported	Deaths, all causes
	Cases, esti- mated expect- ancy	Cases re- ported	Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		Cases, esti- mated expect- ancy	Cases re- ported	Deaths re- ported		
WEST SOUTH CENTRAL											
Arkansas:											
Fort Smith.....	1	1	0	0			0	0		2	
Little Rock.....	3	3	0	0	0	0	0	1	0	0	2
Louisiana:											
New Orleans.....	9	9	0	0	0	4	2	5	2	0	122
Shreveport.....	2	2	0	0	0	1	0	0	1	3	30
Oklahoma:											
Muskogee.....		0		0	0	0		0	0	0	
Texas:											
Dallas.....	9	10	1	0	0	5	0	0	0	8	64
Fort Worth.....	2	8	0	1	0	3	0	1	1	0	36
Galveston.....	0	0	0	0	0	1	0	0	0	0	25
Houston.....	3	5	1	1	0	3	0	1	0	0	73
San Antonio.....	1	2	1	0	0	7	0	1	0	0	72
MOUNTAIN											
Montana:											
Billings.....	1	0	0	0	0	0	0	2	0	2	3
Great Falls.....	2	2	0	0	0	0	0	0	0	0	7
Helena.....	1	0	0	0	0	0	0	0	0	1	7
Missoula.....	1	1	0	0	0	0	0	0	0	0	10
Idaho:											
Boise.....	1	0	0	0	0	0	0	0	0	0	8
Colorado:											
Denver.....	13	19	0	0	0	5	0	0	0	7	80
Pueblo.....	1	1	0	0	0	0	0	1	0	4	7
New Mexico:											
Albuquerque.....	1	1	0	0	0	3	0	4	0	0	11
Arizona:											
Phoenix.....	1	0	0	0	0	2	0	0	0	0	
Utah:											
Salt Lake City.....	2	2	1	0	0	0	0	0	0	0	31
Nevada:											
Reno.....	0	0	0	0	0	0	0	0	0	0	8
PACIFIC											
Washington:											
Seattle.....	10	15	1	0			1	0		8	
Spokane.....	8	0	3	2			0	1		4	
Tacoma.....	4	4	1	0	0	1	0	0	0	3	33
Oregon:											
Portland.....	8	1	3	1	0	1	1	0	0	1	74
Salem.....	1	0	0	0	0	0	0	0	0	1	
California:											
Los Angeles.....	27	27	1	0	0	11	1	2	0	18	263
Sacramento.....	3	0	1	0	0	2	0	1	0	0	32
San Francisco.....	15	5	0	3	0	7	0	1	0	9	155

City reports for week ended December 5, 1931—Continued

Division, State, and city	Meningococcus meningitis		Lethargic encephalitis		Pellagra		Poliomyelitis (infantile paralysis)		
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths
NEW ENGLAND									
Maine:									
Portland.....	0	0	0	0	0	0	0	1	0
Massachusetts:									
Boston.....	0	0	0	0	0	0	2	3	0
Fall River.....	1	1	0	0	0	0	0	0	0
Rhode Island:									
Providence.....	0	0	0	0	0	0	0	1	0
MIDDLE ATLANTIC									
New York:									
New York ¹	7	3	1	0	0	0	2	3	0
Syracuse.....	0	1	0	0	0	0	0	0	0
New Jersey:									
Newark.....	0	0	0	0	0	0	0	1	0
Pennsylvania:									
Philadelphia.....	3	2	1	1	0	0	0	1	0
Pittsburgh.....	3	1	0	0	0	0	0	1	1
EAST NORTH CENTRAL									
Indiana:									
Indianapolis.....	13	3	0	0	0	0	0	0	0
Illinois: ¹									
Chicago.....	2	0	1	0	0	0	1	0	0
Michigan:									
Detroit.....	2	0	0	1	0	0	0	0	0
Wisconsin:									
Milwaukee.....	0	0	0	0	0	0	0	1	0
WEST NORTH CENTRAL									
Minnesota:									
Duluth.....	0	0	0	0	0	0	0	1	0
Minneapolis.....	1	0	0	0	0	0	0	1	0
St. Paul.....	0	0	0	0	0	0	0	2	0
Iowa:									
Des Moines.....	0	0	0	0	0	0	0	1	0
Waterloo.....	1	0	0	0	0	0	0	0	0
Missouri:									
Kansas City.....	0	1	0	0	1	0	0	0	0
SOUTH ATLANTIC									
Maryland:									
Baltimore.....	0	0	0	1	0	0	1	1	0
District of Columbia:									
Washington.....	1	1	0	0	0	0	0	1	0
North Carolina:									
Winston-Salem.....	0	0	0	0	1	0	0	0	0
South Carolina:									
Charleston ¹	0	0	0	0	3	0	0	0	0
Columbia.....	0	0	0	0	0	1	0	0	0
Georgia: ¹									
Savannah ¹	0	0	0	0	3	0	0	0	0
Florida:									
Miami.....	0	0	0	0	2	0	0	0	0
EAST SOUTH CENTRAL									
Kentucky:									
Louisville.....	1	1	0	0	0	0	0	0	0
Tennessee:									
Memphis.....	0	2	0	0	0	1	0	0	0
Nashville.....	1	1	0	0	0	0	0	0	0
Alabama:									
Birmingham.....	0	0	0	0	1	1	0	0	0

¹ Typhus fever, 8 cases: 1 case at New York City, N. Y., 1 case at Springfield, Ill., 3 cases at Charleston, S. C., 1 case at Atlanta, Ga., and 2 cases at Savannah, Ga.

City reports for week ended December 5, 1931—Continued

Division, State, and city	Meningococcus meningitis		Lethargic encephalitis		Pellagra		Polioomyelitis (infantile paralysis)		
	Cases	Deaths	Cases	Deaths	Cases	Deaths	Cases, estimated expectancy	Cases	Deaths
WEST SOUTH CENTRAL									
Louisiana:									
New Orleans.....	0	0	0	0	1	1	1	0	0
Texas:									
Dallas.....	0	0	0	0	1	1	0	0	0
Galveston.....	0	0	0	0	0	2	0	0	0
Houston.....	0	1	0	0	0	1	0	1	0
MOUNTAIN									
Utah:									
Salt Lake City.....	1	2	0	0	0	0	0	0	0
PACIFIC									
Washington:									
Seattle.....	3	0	0	0	0	0	0	0	0
Spokane.....	0	0	0	0	0	0	0	1	0
Tacoma.....	0	0	0	0	0	0	0	1	0
California:									
Los Angeles.....	1	0	0	0	1	0	1	0	0
San Francisco.....	2	2	0	0	0	0	1	3	0

The following table gives the rates per 100,000 population for 98 cities for the 5-week period ended December 5, 1931, compared with those for a like period ended December 6, 1930. The population figures used in computing the rates are estimated mid-year populations for 1930 and 1931, respectively, derived from the 1930 census. The 98 cities reporting cases have an estimated aggregate population of more than 33,000,000. The 91 cities reporting deaths have more than 31,500,000 estimated population.

Summary of weekly reports from cities, November 1 to December 5, 1931—Annual rates per 100,000 population compared with rates for the corresponding period of 1930¹

DIPHTHERIA CASE RATES

	Week ended—									
	Nov. 7, 1931	Nov. 8, 1930	Nov. 14, 1931	Nov. 15, 1930	Nov. 21, 1931	Nov. 22, 1930	Nov. 28, 1931	Nov. 29, 1930	Dec. 5, 1931	Dec. 6, 1930
98 cities.....	94	82	96	89	96	100	85	87	101	90
New England.....	84	85	50	82	70	123	67	67	58	121
Middle Atlantic.....	32	33	52	44	53	52	58	48	54	58
East North Central.....	97	109	80	128	91	124	72	122	94	112
West North Central.....	155	77	184	107	174	110	151	110	222	101
South Atlantic.....	182	86	146	120	172	154	144	66	159	112
East South Central.....	268	215	227	185	169	275	145	138	163	143
West South Central.....	203	199	233	160	206	171	207	153	244	147
Mountain.....	44	123	61	26	17	26	27	79	52	18
Pacific.....	100	93	127	63	98	63	67	95	88	65

MEASLES CASE RATES

98 cities.....	44	59	55	91	85	126	91	107	114	143
New England.....	161	128	238	172	233	179	315	162	481	220
Middle Atlantic.....	27	34	38	68	92	76	82	69	111	85
East North Central.....	18	16	18	17	29	31	15	28	31	28
West North Central.....	15	282	17	502	19	767	15	649	27	953
South Atlantic.....	12	48	10	26	34	64	28	44	44	62
East South Central.....	12	84	12	18	29	149	35	66	35	155
West South Central.....	27	0	24	0	10	3	24	10	27	11
Mountain.....	444	229	400	308	757	326	1,277	282	757	53
Pacific.....	104	24	135	32	149	28	123	10	180	26

SCARLET FEVER CASE RATES

98 cities.....	169	169	170	187	187	195	156	174	179	202
New England.....	202	225	221	276	260	237	262	264	293	298
Middle Atlantic.....	134	133	131	126	163	159	147	148	155	178
East North Central.....	239	231	215	287	241	263	171	221	229	257
West North Central.....	140	140	149	143	132	219	123	139	161	198
South Atlantic.....	190	158	239	154	259	216	176	188	175	230
East South Central.....	99	293	198	275	145	209	122	215	128	299
West South Central.....	95	91	122	118	78	94	93	132	108	92
Mountain.....	252	282	313	388	218	282	198	229	218	141
Pacific.....	121	95	96	99	129	87	108	83	100	97

SMALLPOX CASE RATES

98 cities.....	3	2	1	4	1	3	3	8	6	7
New England.....	0	0	0	0	0	0	0	0	55	0
Middle Atlantic.....	0	0	0	0	0	0	0	0	1	0
East North Central.....	0	4	0	2	0	0	0	4	0	1
West North Central.....	11	6	4	21	10	23	13	68	4	48
South Atlantic.....	0	0	0	0	0	0	0	0	0	0
East South Central.....	12	0	6	0	0	0	6	0	0	0
West South Central.....	3	7	3	3	0	3	21	3	3	4
Mountain.....	0	9	9	0	0	44	0	35	0	108
Pacific.....	6	6	4	18	6	6	6	8	10	10

¹ The figures given in this table are rates per 100,000 population, annual basis, and not the number of cases reported. Populations used are estimated as of July 1, 1931, and 1930, respectively.

² Waterloo, Iowa, not included.

³ South Bend, Ind., St. Paul, Minn., Fort Smith, Ark., and Reno, Nev., not included.

⁴ Lynchburg, Va., not included.

⁵ Shreveport, La., not included.

⁶ South Bend, Ind., not included.

⁷ St. Paul, Minn., not included.

⁸ Fort Smith, Ark., not included.

⁹ Reno, Nev., not included.

Summary of weekly reports from cities, November 1 to December 5, 1931—Annual rates per 100,000 population compared with rates for the corresponding period of 1930—Continued

TYPHOID FEVER CASE RATES

	Week ended—									
	Nov. 7, 1931	Nov. 8, 1930	Nov. 14, 1931	Nov. 15, 1930	Nov. 21, 1931	Nov. 22, 1930	Nov. 28, 1931	Nov. 29, 1930	Dec. 5, 1931	Dec. 6, 1930
98 cities.....	12	11	12	15	12	15	17	10	17	10
New England.....	10	5	7	24	10	17	2	12	5	7
Middle Atlantic.....	11	5	6	4	8	5	4	3	5	8
East North Central.....	6	9	11	5	5	9	16	4	4	10
West North Central.....	21	14	13	19	6	23	19	8	4	6
South Atlantic.....	30	32	36	34	24	28	34	32	16	18
East South Central.....	17	24	23	48	41	12	6	12	12	12
West South Central.....	30	28	24	87	41	84	17	70	27	26
Mountain.....	9	18	0	26	9	53	10	9	26	9
Pacific.....	0	16	10	10	18	10	2	6	10	10

INFLUENZA DEATH RATES

	7		8		7		10		17		9		17		19	
	7	9	8	9	7	10	17	9	17	9	17	19	17	19	17	19
91 cities.....	12	2	14	5	7	7	0	2	2	2	2	2	2	2	2	2
New England.....	12	2	14	5	7	7	0	2	2	2	2	2	2	2	2	2
Middle Atlantic.....	8	12	10	8	6	7	9	11	4	4	4	4	4	4	4	4
East North Central.....	5	6	2	9	4	5	5	7	6	6	6	6	6	6	6	6
West North Central.....	6	3	6	6	6	6	3	0	6	6	6	6	6	6	6	6
South Atlantic.....	4	10	6	6	12	24	6	10	14	14	14	14	14	14	14	14
East South Central.....	0	26	0	39	25	13	13	26	38	38	38	38	38	38	38	38
West South Central.....	17	14	7	28	10	36	17	14	7	7	7	7	7	7	7	7
Mountain.....	17	9	27	9	17	62	27	26	9	9	9	9	9	9	9	9
Pacific.....	5	7	12	5	5	7	7	7	19	19	19	19	19	19	19	19

PNEUMONIA DEATH RATES

	88		101		101		116		186		109		189		199	
	88	101	88	101	101	116	186	109	189	199	189	199	189	199	189	199
91 cities.....	67	89	101	114	84	126	99	77	91	73	91	73	91	73	91	73
New England.....	67	89	101	114	84	126	99	77	91	73	91	73	91	73	91	73
Middle Atlantic.....	107	116	106	129	116	133	98	118	95	101	95	101	95	101	95	101
East North Central.....	64	74	52	85	70	82	52	78	56	77	56	77	56	77	56	77
West North Central.....	80	87	88	78	115	138	119	93	88	132	88	132	88	132	88	132
South Atlantic.....	117	152	97	172	152	186	122	180	146	154	146	154	146	154	146	154
East South Central.....	129	136	151	188	183	175	107	136	95	155	95	155	95	155	95	155
West South Central.....	66	110	55	103	79	114	66	153	135	128	135	128	135	128	135	128
Mountain.....	139	194	148	220	174	167	126	229	122	132	122	132	122	132	122	132
Pacific.....	53	42	70	67	50	50	74	70	77	60	77	60	77	60	77	60

¹ Waterloo, Iowa, not included.

² South Bend, Ind., St. Paul, Minn., Fort Smith, Ark., and Reno, Nev., not included.

³ Lynchburg, Va., not included.

⁴ Shreveport, La., not included.

⁵ South Bend, Ind., not included.

⁶ St. Paul, Minn., not included.

⁷ Fort Smith, Ark., not included.

⁸ Reno, Nev., not included.

⁹ South Bend, Ind., St. Paul, Minn., and Reno, Nev., not included.

FOREIGN AND INSULAR

BRITISH GUIANA

Deaths from certain diseases—1928, 1929, 1930.—According to the annual report of the Surgeon General of British Guiana for the year 1930, deaths from certain diseases were reported in the colony during the years 1928, 1929, and 1930, as follows:

Disease	1928	1929	1930	Disease	1928	1929	1930
Ancylostomiasis.....	33	10	28	Influenza.....	91	121	94
Blackwater fever.....	6	11	12	Malaria.....	1,563	1,198	1,104
Diarrhea and enteritis.....	557	448	380	Nephritis.....	694	514	528
Dysentery.....	185	141	105	Pneumonia.....	711	661	588
Filariasis.....	47	52	37	Tuberculosis.....	301	276	302
Heart disease.....	363	351	359	Typhoid fever.....	58	44	53

Population Dec. 31, 1930, 312,489.

CANADA

Provinces—Communicable diseases—Week ended November 28, 1931.—The Department of Pensions and National Health of Canada reports cases of certain communicable diseases for the week ended November 28, 1931, as follows:

Province	Cerebro-spinal fever	Influenza	Lethargic encephalitis	Polio-myelitis	Smallpox	Typhoid fever
Prince Edward Island ¹						
Nova Scotia.....		19			1	3
New Brunswick.....						4
Quebec.....	2			17		15
Ontario.....	1	6	2		2	11
Manitoba.....	1				1	8
Saskatchewan.....					5	
Alberta.....					2	
British Columbia.....						1
Total.....	4	25	2	17	11	42

¹ No case of any disease included in the table was reported during the week.

Quebec Province—Communicable diseases—Week ended November 28, 1931.—The Bureau of Health of the Province of Quebec, Canada, reports cases of certain communicable diseases for the week ended November 28, 1931, as follows:

Disease	Cases	Disease	Cases
Cerebrospinal meningitis.....	2	Paratyphoid fever.....	1
Chicken pox.....	142	Polio-myelitis.....	17
Diphtheria.....	53	Scarlet fever.....	50
Erysipelas.....	6	Tuberculosis.....	17
German measles.....	9	Typhoid fever.....	14
Measles.....	165	Whooping cough.....	73
Mumps.....	29		

CUBA

Habana—Communicable diseases—Four weeks ended November 7, 1931.—During the four weeks ended November 7, 1931, certain communicable diseases were reported in the city of Habana, Cuba, as follows:

Disease	Cases	Deaths	Disease	Cases	Deaths
Chicken pox.....	2		Poliomyelitis.....	2	
Diphtheria.....	9	1	Scarlet fever.....	2	
Leprosy.....	2		Tuberculosis.....	23	4
Malaria.....	18		Typhoid fever.....	9	4
Measles.....	54				

GREAT BRITAIN

England and Wales—Vital statistics—July–September, 1931.—During the third quarter of the year 1931, 161,267 births and 96,745 deaths were registered in England and Wales, giving a birth rate on an annual basis of 16.0 per 1,000 population and a death rate of 9.6 per 1,000. The figures are provisional. The mortality of infants under 1 year of age was 45 per 1,000 live births.

During the 13 weeks ended October 3, 1931, deaths from certain communicable diseases were reported in 107 boroughs and great towns, including Greater London, as follows:

Disease	Number of deaths	Death rate per 1,000 population	Disease	Number of deaths	Death rate per 1,000 population
Diarrhea and enteritis (under 2 years).....	566		Scarlet fever.....	44	0.01
Diphtheria.....	298	0.06	Smallpox.....	0	
Influenza.....	259	.05	Typhoid fever.....	18	
Measles.....	146	.03	Whooping cough.....	315	.03

Deaths from certain communicable diseases in 159 smaller towns for the quarter ended September 30, 1931, were as follows:

Disease	Deaths	Disease	Deaths
Diarrhea and enteritis (under 2 years).....	73	Scarlet fever.....	10
Diphtheria.....	46	Smallpox.....	0
Influenza.....	63	Typhoid fever.....	9
Measles.....	35	Whooping cough.....	23

England and Wales—Communicable diseases—Thirteen weeks ended October 3, 1931.—During the 13 weeks ended October 3, 1931, cases of certain communicable diseases were reported in England and Wales as follows (civilians only):

Disease	Cases	Disease	Cases
Diphtheria.....	10,820	Puerperal pyrexia.....	1,406
Ophthalmia neonatorum.....	1,369	Scarlet fever.....	18,941
Pneumonia.....	6,701	Smallpox.....	459
Puerperal fever.....	512	Typhoid fever.....	737

Scotland—Vital statistics—Quarter ended September 30, 1931.—The Registrar General of Scotland has published the following statistics for the third quarter of the year 1931:

Population (provisional).....	4,842,554	Deaths from—Continued.	
Births.....	22,659	Heart disease.....	1,016
Birth rate per 1,000 population.....	18.6	Influenza.....	56
Deaths.....	13,242	Pneumonia.....	133
Death rate per 1,000 population.....	10.8	Pneumonia, lobar.....	185
Marriages.....	9,351	Measles.....	97
Deaths under 1 year.....	1,353	Nephritis (acute).....	46
Deaths under 1 year per 1,000 births.....	60	Nephritis (chronic).....	259
Deaths from—		Puerperal sepsis.....	31
Bronchitis.....	433	Scarlet fever.....	19
Broncho-pneumonia.....	304	Syphilis.....	24
Cerebrospinal fever.....	54	Tetanus.....	2
Diabetes.....	164	Tuberculosis.....	928
Diphtheria.....	69	Typhoid fever.....	6
Dysentery.....	2	Whooping cough.....	121
Erysipelas.....	29		

SWITZERLAND

Deaths from tuberculosis—1911–1920, 1921–1930.—According to a recent report, deaths from all forms of tuberculosis occurred in Switzerland, during the 10-year periods 1911–1920 and 1921–1930, as follows:

Age group	Deaths			
	1911–1920		1921–1930	
	Males	Females	Males	Females
0–14.....	4,757	5,505	2,588	2,988
15–29.....	9,495	13,911	7,039	11,353
30–49.....	12,276	11,459	8,737	8,120
50–69.....	8,148	7,115	7,135	5,962
70 and over.....	1,488	2,241	1,406	2,158
Total.....	36,164	40,231	26,905	30,601

The population of Switzerland, according to the census of Dec. 31, 1930, is 4,082,511.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

CHOLERA—Continued

[C indicates cases; D, deaths; P, present]

Place	Week ended—																				
	May 31- June 27, 1931	June 28- July 25, 1931	July 26- Aug. 22, 1931	Aug. 23- Sept. 19, 1931	October, 1931					November, 1931			December, 1931								
					3	10	17	24	31	7	14	21	28	5	12						
Philippine Islands: 1																					
Provinces—																					
Cebu.....	4	4	3	35	49	21	5	4							7	5			4	16	7
Iloilo.....	20	27	3	16	35	16	5	3							5	4			4	10	5
Negros Occidental.....	2	20																			
Slam.....	1	4	1																		
Bangkok.....	2	1	1	1																	
On vessel:	1																				
S. S. City of Eastborne, at Calcutta, from Cocosnada.....	1																				
S. S. Tairea, at Penang, from Calcutta.....																					
S. S. Bandar Shalpour, at Bushire, Persia, from Basra.....			1																		
S. S. Kohistan, at Basra, from Bushire, Persia.....			2																		
S. S. Cathay, at Kobe, Japan, from Shanghai.....			4																		
S. S. Kasagi Maru, at Moli, from Shanghai.....			1	1																	
S. S. Ankoo, at Nagasaki, from Shanghai.....				1																	

¹ Figures for cholera in the Philippine Islands are subject to correction.

Place	May, 1931	June, 1931	July, 1931	August, 1931			September, 1931			October, 1931			Nov. 1-10, 1931
				1-10	11-20	21-31	1-10	11-20	21-30	1-10	11-20	21-31	
Indo-China (French) (see also table above):													
Cambodia ¹	117	308	241	12					8	6	1	16	2
Cochin-China ¹	63	109	60	2					4	3	1	16	1
	174	140	143	39					9	9	11	2	1
	133	106	42	32					6	7	10	2	4

¹ Reports incomplete.

PLAGUE

[C indicates cases; D, deaths; P, present]

Place	May 31- June 27, 1931	June 28- July 26, 1931	July 27- Aug. 25, 1931	Aug. 26- Sept. 24, 1931	Week ended—									
					Sept. 25, 1931			October, 1931			November, 1931			December, 1931
					3	10	17	24	31	7	14	21	28	
Algeria:														
Algiers.....	1	1	2											
Bone.....														
Philippeville.....		P												
Argentina: San Juan Province.....														
Belgian Congo.....	1													
British East Africa (see also table below):														
Tanganyika.....	17	6	8	4	8	2	3							
Uganda.....	10	6	2	4	4									
	298	418	285	280	83	62	64	71						
Ceylon: Colombo.....	286	400	281	207	82	63	58	69						
	2	1	6	3	1	1	1							
Plague-infected rats.....	2	1	6	3	1	1	1							
Chile: Santiago.....							1							

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

PLAGUE—Continued

[O indicates cases; D, deaths; P, present]

Place	June, 1931	July, 1931	August, 1931	September, 1931	October, 1931	November, 1931	Place	June, 1931	July, 1931	August, 1931	September, 1931	October, 1931	November, 1931
British East Africa (see also table above):							Madagascar—Continued.						
Kenya.....	O						Moromanga Province.....	O					
Ecuador:							Tananarive Province.....	O					
Alamora Parish—Los Hoyos.....	O						Peru.....	O					
Amaluza Parish—Cangochapa.....	O						Senegal:						
Calvas Canton—							Baol.....	O					
Carlmanaga.....	O						Dakar.....	O					
Ovejeria.....	O						Diourbel.....	O					
Colcia Canton—Oboras.....	O						Louga.....	O					
Loja Canton—	O						Rufisque.....	O					
Lapaz.....	O						Thies.....	O					
Naimuro.....	O						Tivouane.....	O					
Paterillo.....	O												
Tuburo.....	O												
Palas Canton—San Antonio.....	O												
Indo-China (see also table above):													
Madagascar (see also table above):													
Ambositra Province.....	O												
Antsirabe Province.....	O												
Marianarivo Province.....	O												

! Reports incomplete.

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

SMALLPOX—Continued

[C indicates cases; D, death; P, present]

[illegible]

TYPHUS FEVER

Place	Week ended—																		
	May 31— June 27, 1931	June 28— July 25, 1931	July 26— Aug. 22, 1931	September, 1931					October, 1931				November, 1931						
				Aug. 29, 1931	5	12	19	26	3	10	17	24	31	7	14	21	28		
Algeria:																			
Algiers	9	2	2																
Bone	26	3	1																
Constantine Department	2		1																
Oran	30			1															
Bulgaria	6																		
Chile:																			
Antofagasta																			
Santiago																			
China:																			
Manchuria—Harbin	16	3	1																
Shanghai																			
Chosen (see table below)																			
Colombia: Cali																			
Czechoslovakia (see table below)																			
Egypt:																			
Alexandria	3																		
Beheira																			
Cairo																			
Gharbieh																			
Greece (see table below)																			
Guatemala (see table below)																			
Irish Free State:																			
Cork County—																			
Schull		1																	
Skibbereen		1																	
Kerry County—Listowel																			
Limerick County—																			
Croom																			
Glin																			
Limerick		2	1																
Michelstown		1																	
Rathkeale																			
Mayo County—																			
Castlebar			1																
Westport			1																
Japan																			

CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

YELLOW FEVER—Continued

(C indicates cases; D, deaths; P, present)

[illegible]

Ad

Air
Ala

Alc
Anc
in
Ant
Ant
Arg
Ari

Ar

Ba

Ba
i
Bir
s
Br
Br
Br
Br

Ca

Ca

INDEX

A

Admissions to hospitals for the insane:	Page
July, 1929.....	3073
August, 1929.....	3124
Air jet hydrocyanic acid sprayer—Williams.....	1755
Alabama:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Alcaligenes faecalis, a new subspecies of, radicans—Evans.....	1676
Anopheles atropos D. and K.—A new potential carrier of malaria organisms—Mayne and Griffiths.....	3107
Anthrax. (See Summaries—Disease cases reported monthly by States.)	
Antifreeze mixtures, Arkansas law relating to.....	2254
Argentina: San Juan Province—Plague (human).....	1894
Arizona:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Arkansas:	
(See also Name of disease—Current weekly State reports.)	
(See also Summaries—Disease cases reported monthly by States.)	
Law relating to antifreeze mixtures.....	2254

B

Badger, L. F.:	
Experimental transmission of endemic typhus fever of the United States by the rat flea (<i>Xenopsylla cheopis</i>).....	2415
Leprosy. (A study of white blood cells and their relation to clinical progress).....	2782
Typhus fever. (The experimental transmission of endemic typhus fever of the United States by the rat flea <i>Xenopsylla cheopis</i>).....	2481
Typhus fever. (The rat flea, <i>Xenopsylla cheopis</i> , in experimental transmission).....	1869
Typhus fever. Typhus virus in feces of infected fleas (<i>Xenopsylla cheopis</i>) and duration of infectivity of fleas.....	3103
Baker, C. E.: Microscopic examination for intestinal parasites of 73 boys in the National Training School for Boys, Washington, D. C.....	2980
Birth, death, and infant mortality figures, provisional, birth registration area, 1930.....	2373
British Cameroons: Mamfe—Yellow fever, May 28, 1931.....	1656
British Guiana: Deaths from certain diseases, 1928, 1929, and 1930.....	3137
Britten, Rollo H.: Physical examination as an instrument of research.....	1671
Brundage, Dean K.:	
A survey of the work of employees' mutual benefit associations.....	2102
Sickness among male industrial employees in the first quarter of 1931.....	1799
Sickness among male industrial employees in the second quarter of 1931.....	2499

C

California:	
(See also Name of disease—Current weekly State reports.)	
(See also Summaries—Disease cases reported monthly by States.)	
San Benito County—Plague-infected ground squirrels.....	1954, 2133
Canada:	
Ontario—Communicable diseases—Four weeks ended—	
June 27, 1931.....	1854
July 25, 1931.....	2143
August 29, 1931 (five weeks).....	2396
October 31, 1931 (five weeks).....	3084

Canada—Continued.

Provinces—Communicable diseases—Week ended—	Page
June 6, 1931.....	1599
June 13, 1931.....	1656
June 20, 1931.....	1711
June 27, 1931.....	1784
July 4, 1931.....	1853
July 11, 1931.....	1894
July 18, 1931.....	1965
July 25, 1931.....	2026
August 1, 1931.....	2077
August 8, 1931.....	2143
August 15, 1931.....	2205
August 22, 1931.....	2273
August 29, 1931.....	2339
September 5, 1931.....	2395
September 12, 1931.....	2463
September 19, 1931.....	2521
September 26, 1931.....	2645
October 3, 1931.....	2645
October 10, 1931.....	2691
October 17, 1931.....	2757
October 24, 1931.....	2821
October 31, 1931.....	2880
November 7, 1931.....	2956
November 14, 1931.....	3006
November 21, 1931.....	3084
November 28, 1931.....	3137
Quebec—Communicable diseases—Week ended—	
June 13, 1931.....	1599
June 20, 1931.....	1656
June 27, 1931.....	1711
July 4, 1931.....	1784
July 11, 1931.....	1853
July 18, 1931.....	1894
July 25, 1931.....	1965
August 1, 1931.....	2026
August 8, 1931.....	2077
August 15, 1931.....	2205
August 22, 1931.....	2273
August 29, 1931.....	2339
September 5, 1931.....	2395
September 12, 1931.....	2463
September 19, 1931.....	2521
September 26, 1931.....	2582
October 3, 1931.....	2645
October 10, 1931.....	2757
October 17, 1931.....	2691
October 24, 1931.....	2821
October 31, 1931.....	2880
November 7, 1931.....	2956
November 14, 1931.....	3006
November 21, 1931.....	3084
November 28, 1931.....	3137
Saskatchewan—Vital statistics—1930.....	3006
Carnes, E. H.: Rat-flea survey of the port of St. Thomas, Virgin Islands.....	2558
Case, Jarvis D.: A study of illness among grade-school children.....	1801
Catalytic action of copper in the oxidation of crystalline glutathione— Voegtlin, Johnson, and Rosenthal.....	2234
Ceder, E. T.: Experimental transmission of endemic typhus fever of the United States by the rat flea (<i>Xenopsylla cheopis</i>).....	2415
Typhus fever. (The experimental transmission of endemic typhus of the United States by the rat flea, <i>Xenopsylla cheopis</i>).....	2481
Typhus fever. (The rat flea, <i>Xenopsylla cheopis</i> , in experimental transmission).....	1869
Typhus fever. Typhus virus in feces of infected fleas (<i>Xenopsylla cheopis</i>) and duration of infectivity of fleas.....	3103

Cell division, chemistry of. II. The relation between cell growth and division in <i>Amoeba proteus</i> —Chalkley.....	1736
Chalkley, H. W.: Chemistry of cell division. II. The relation between cell growth and division in <i>Amoeba proteus</i>	1736
Chemistry of cell division. II. The relation between cell growth and division in <i>Amoeba proteus</i> —Chalkley.....	1736
Chicken pox: (See also Summaries—Disease cases reported monthly by States.) City reports for week ended—	
June 13, 1931.....	1590
June 20, 1931.....	1647
June 27, 1931.....	1702
July 4, 1931.....	1774
July 11, 1931.....	1844
July 18, 1931.....	1885
July 25, 1931.....	1955
August 1, 1931.....	2017
August 8, 1931.....	2067
August 15, 1931.....	2134
August 22, 1931.....	2195
August 29, 1931.....	2264
September 5, 1931.....	2329
September 12, 1931.....	2385
September 19, 1931.....	2453
September 26, 1931.....	2511
October 3, 1931.....	2572
October 10, 1931.....	2635
October 17, 1931.....	2681
October 24, 1931.....	2748
October 31, 1931.....	2812
November 7, 1931.....	2870
November 14, 1931.....	2946
November 21, 1931.....	2996
November 28, 1931.....	3075
December 5, 1931.....	3127
Child, health of the school.....	2605
Children, a study of illness among grade-school—Watkins, Wilson, Hiseck, Case, and Rice.....	1801
China: Changchow—Plague (human).....	1894
Chiobe—Plague (human).....	1894
Manchuria—Fumigation of vessels at Dairen and Port Arthur.....	2144
Shanghai—Meningococcus meningitis.....	1711
Shansi Province— Plague (human).....	2646, 2821
Vital statistics—Year, 1923.....	2582
Shensi Province—Plague (human).....	2821
Cholera: Foreign reports.....	1601,
1658, 1713, 1786, 1856, 1896, 1967, 2027, 2078, 2145, 2208,	
2276, 2341, 2399, 2465, 2523, 2585, 2647, 2693, 2759, 2823,	
2882, 2958, 3008, 3087, 3140.	
Iraq.....	3007
Basra.....	1966
Persia.....	2692
City health officers, 1931, directory of.....	2921
Clinical tropical medicine, special course in.....	1640
Cockroaches, experiments with certain fumigants used for the destruction of—Ridlon.....	1623
Cockroaches on ships, effect of fumigation on—Williams.....	1680
Collins, Selwyn D.: Age and sex incidence of influenza and pneumonia morbidity and mortality in the epidemic of 1928-29 with comparative data for the epidemic of 1918-19.....	1909

Colorado:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Communicable diseases:

Canada—

Ontario—Four weeks ended—

	Page
June 27, 1931.....	1854
July 25, 1931.....	2143
August 29, 1931 (five weeks).....	2396
October 31, 1931 (five weeks).....	3084

Provinces—Week ended—

June 6, 1931.....	1599
June 13, 1931.....	1656
June 20, 1931.....	1711
June 27, 1931.....	1784
July 4, 1931.....	1853
July 11, 1931.....	1894
July 18, 1931.....	1965
July 25, 1931.....	2026
August 1, 1931.....	2077
August 8, 1931.....	2143
August 15, 1931.....	2205
August 22, 1931.....	2273
August 29, 1931.....	2339
September 5, 1931.....	2395
September 12, 1931.....	2463
September 19, 1931.....	2521
September 26, 1931.....	2645
October 3, 1931.....	2691
October 10, 1931.....	2757
October 17, 1931.....	2821
October 24, 1931.....	2880
October 31, 1931.....	2956
November 7, 1931.....	3006
November 14, 1931.....	3084
November 21, 1931.....	3137
November 28, 1931.....	

Quebec—Week ended—

June 13, 1931.....	1599
June 20, 1931.....	1656
June 27, 1931.....	1711
July 4, 1931.....	1784
July 11, 1931.....	1853
July 18, 1931.....	1894
July 25, 1931.....	1965
August 1, 1931.....	2026
August 8, 1931.....	2077
August 15, 1931.....	2205
August 22, 1931.....	2273
August 29, 1931.....	2339
September 5, 1931.....	2395
September 12, 1931.....	2463
September 19, 1931.....	2521
September 26, 1931.....	2582
October 3, 1931.....	2645
October 10, 1931.....	2757
October 17, 1931.....	2691
October 24, 1931.....	2821
October 31, 1931.....	2880
November 7, 1931.....	2956
November 14, 1931.....	3006
November 21, 1931.....	3084
November 28, 1931.....	3137

Costa Rica—San Jose—January–April, 1931.....

1784

Communicable diseases—Continued.

Cuba—

Habana—

	Page
June 20, 1931 (four weeks)	1712
July 18, 1931 (four weeks)	2205
August 15, 1931 (four weeks)	2273
September 12, 1931 (four weeks)	2463
October 10, 1931 (four weeks)	3085
November 7, 1931 (four weeks)	3138

Provinces—

June 6, 1931 (four weeks)	1657
July 4, 1931 (three weeks)	2274
August 1, 1931 (four weeks)	2339
August 29, 1931 (four weeks)	2583
September 26, 1931 (four weeks)	3085

Czechoslovakia—

April, 1931	1599
May, 1931	1895
June, 1931	2464
July, 1931	2522
August, 1931	2691
September, 1931	3006

Denmark—

April, 1931	1712
May, 1931	2144
June, 1931	2396
July, 1931	2522
August, 1931	2583
September, 1931	3086

Foreign reports

1658, 1713, 1786, 1856, 1896, 1967, 2027, 2078, 2145, 2208, 2276, 2341, 2399, 2465, 2523, 2585, 2647, 2693, 2759, 2823, 2882, 2958, 3008, 3087, 3140.	1601,
---	-------

Great Britain—England and Wales—

Quarter ended July 4, 1931	2397
Quarter ended October 3, 1931	3139

Jamaica—Four weeks ended—

June 20, 1931	1785
July 18, 1931	2206
August 15, 1931	2274
September 12, 1931	2464
October 10, 1931	2692

Latvia—

January-June, 1931	2340
July, 1931	2522
August, 1931	2822
September, 1931	2880

Mexico—Tampico—

June, 1931	1854
July, 1931	2206
August, 1931	2340
September, 1931	2692
October, 1931	2881

Panama Canal Zone—

May, 1931	1855
June, 1931	1966
July, 1931	2275
August, 1931	2583
September, 1931	2757
October, 1931	3086

Porto Rico—San Juan—Four weeks ended—

June 20, 1931	1855
July 18, 1931	2206
August 15, 1931	2398
September 12, 1931	2522
October 10, 1931	2758
November 7, 1931	2881

Communicable diseases—Continued.

United States—

Case rates per 100,000 population for 98 cities—

	Page
May 10–June 13, 1931.....	1597
May 17–June 20, 1931.....	1654
May 24–June 27, 1931.....	1709
May 31–July 4, 1931.....	1782
June 7–July 11, 1931.....	1851
June 14–July 18, 1931.....	1892
June 21–July 25, 1931.....	1962
June 28–August 1, 1931.....	2024
July 5–August 8, 1931.....	2075
July 12–August 15, 1931.....	2141
July 19–August 22, 1931.....	2202
July 26–August 29, 1931.....	2271
August 2–September 5, 1931.....	2337
August 9–September 12, 1931.....	2393
August 16–September 19, 1931.....	2461
August 23–September 26, 1931.....	2519
August 30–October 3, 1931.....	2580
September 6–October 10, 1931.....	2643
September 13–October 17, 1931.....	2688
September 20–October 24, 1931.....	2755
September 27–October 31, 1931.....	2819
October 4–November 7, 1931.....	2878
October 11–November 14, 1931.....	2954
October 18–November 21, 1931.....	3004
October 25–November 28, 1931.....	3082
November 1–December 5, 1931.....	3135

Cases and case rates per 100,000 population—Reported by State health officers—

February, 1931.....	1771, 1772
March, 1931.....	2451, 2452
April, 1931.....	2509, 2510
May, 1931.....	2569, 2570
June, 1931.....	2679, 2680
July, 1931.....	2810, 2811
August, 1931.....	2993, 2994
September, 1931.....	3071, 3072

Current prevalence of..... 1615, 1871, 2093, 2357, 2601, 2837, 3115

Current weekly State reports..... 1585,

1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129, 2191, 2259,
2324, 2380, 2447, 2506, 2566, 2629, 2675, 2744, 2806, 2866, 2940,
2989, 3067, 3121.

Reciprocal notifications—

April, 1931.....	2194
May, 1931.....	2262
June, 1931.....	2328
July, 1931.....	2384
August, 1931.....	2452
September, 1931.....	2633
October, 1931.....	2945

Summary—Disease cases reported monthly by States..... 1587,

1645, 1699, 1770, 1842, 1882, 1952, 2015, 2065, 2131, 2193, 2261,
2326, 2382, 2449, 2508, 2568, 2631, 2677, 2746, 2808, 2868, 2942,
2991, 3069, 3123.

Virgin Islands—

May, 1931.....	1657
June, 1931.....	1855
July, 1931.....	2077
August, 1931.....	2398
September, 1931.....	2646
October, 1931.....	2957

Yugoslavia—

May, 1931.....	1600
June, 1931.....	2207
July, 1931.....	2275

Communicable diseases—Continued.

Yugoslavia—Continued.	Page
August, 1931.....	2398
September, 1931.....	2822
October, 1931.....	3007
Conjunctivitis. (See Summaries—Disease cases reported monthly by States.)	
Connecticut:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Copper, catalytic action of, in the oxidation of crystalline glutathione—	
Voegtlin, Johnson, and Rosenthal.....	2234
Costa Rica: San Jose—Communicable diseases—January–April, 1931.....	1784
Court decisions—Relating to public health.....	1582,
1694, 1763, 1836, 1876, 1946, 2008, 2060, 2125, 2187, 2320, 2376,	
2443, 2502, 2562, 2625, 2740, 2802, 2984, 3064, 3117.	
Cuba:	
Habana—Communicable diseases—	
June 20, 1931 (four weeks).....	1712
July 18, 1931 (four weeks).....	2205
August 15, 1931 (four weeks).....	2273
September 12, 1931 (four weeks).....	2463
October 10, 1931 (four weeks).....	3085
November 7, 1931 (four weeks).....	3138
Provinces—Communicable diseases—	
June 6, 1931 (four weeks).....	1657
July 4, 1931 (three weeks).....	2274
August 1, 1931 (four weeks).....	2339
August 29, 1931 (four weeks).....	2583
September 26, 1931 (four weeks).....	3085
Cultures, hanging drop tissue, a technique for adjustment of pH of—	
Earle.....	1998
Cumming, Hugh S.: Present-day problems of yellow fever.....	2361
Current weekly State reports. (See Name of disease—Current weekly State reports.)	
Cyanogen product in ship fumigation, report on some tests of the use of a new—Williams.....	2048
Czechoslovakia: Communicable diseases—	
April, 1931.....	1599
May, 1931.....	1895
June, 1931.....	2464
July, 1931.....	2522
August, 1931.....	2691
September, 1931.....	3006

D

Death, birth, and infant mortality figures, provisional, birth registration area, 1930.....	2373
Death rates—Group of insured persons—Metropolitan Life Insurance Co. compilation—	
May, 1931.....	1762
June, 1931.....	2059
July, 1931.....	2253
August, 1931.....	2623
September, 1931.....	2860
Deaths. (See also Mortality.)	
All causes—City reports for week ended—	
June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020

Deaths—Continued.

All causes—City reports for week ended—Continued.	Page
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750
October 31, 1931.....	2815
November 7, 1931.....	2873
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130
British Guiana, 1928, 1929, and 1930.....	3137
Ecuador—Guayaquil—1930.....	2026
Large cities—Week ended—	
June 13, 1931.....	1582
June 20, 1931.....	1640
June 27, 1931.....	1694
July 4, 1931.....	1765
July 11, 1931.....	1837
July 18, 1931.....	1877
July 25, 1931.....	1947
August 1, 1931.....	2010
August 8, 1931.....	2060
August 15, 1931.....	2126
August 22, 1931.....	2188
August 29, 1931.....	2256
September 5, 1931.....	2321
September 12, 1931.....	2377
September 19, 1931.....	2444
September 26, 1931.....	2503
October 3, 1931.....	2563
October 10, 1931.....	2626
October 17, 1931.....	2672
October 24, 1931.....	2741
October 31, 1931.....	2803
November 7, 1931.....	2863
November 14, 1931.....	2937
November 21, 1931.....	2986
November 28, 1931.....	3064
December 5, 1931.....	3118
Mexico—June 1-28, 1931.....	1785
Delaware:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Dengue. (See Summaries—Disease cases reported monthly by States.)	
Denmark: Communicable diseases—	
April, 1931.....	1712
May, 1931.....	2144
June, 1931.....	2396
July, 1931.....	2522
August, 1931.....	2583
September, 1931.....	3086
Dental decay and corrections among school children of different ages—	
Stoughton and Meaker.....	2608
Dermatitis venenata due to contact with Brazilian walnut wood—	
Schwartz.....	1938
Diesel motor boats, rat population on.....	2371

Diphtheria:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—

	Page
June 13, 1931.....	1590
June 20, 1931.....	1647
June 27, 1931.....	1702
July 4, 1931.....	1774
July 11, 1931.....	1844
July 18, 1931.....	1885
July 25, 1931.....	1955
August 1, 1931.....	2017
August 8, 1931.....	2067
August 15, 1931.....	2134
August 22, 1931.....	2195
August 29, 1931.....	2264
September 5, 1931.....	2329
September 12, 1931.....	2385
September 19, 1931.....	2453
September 26, 1931.....	2511
October 3, 1931.....	2572
October 10, 1931.....	2635
October 17, 1931.....	2681
October 24, 1931.....	2748
October 31, 1931.....	2812
November 7, 1931.....	2870
November 14, 1931.....	2946
November 21, 1931.....	2996
November 28, 1931.....	3075
December 5, 1931.....	3127
Current weekly State reports.....	1585,
1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129,	2191,
2259, 2324, 2380, 2447, 2506, 2566, 2629, 2675, 2744,	2806,
2866, 2940, 2989, 3067, 3121.	

Directory:

City health officers, 1931.....	2921
County health officers, 1931.....	3051
State and insular health authorities, 1931.....	2899

District of Columbia:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Draper, W. F.: Some essential considerations in connection with the rural health program.....	1617
---	------

Dyer, R. E.:

Experimental transmission of endemic typhus fever of the United States by the rat flea (<i>Xenopsylla cheopis</i>).....	2415
Typhus fever. (The experimental transmission of endemic typhus fever of the United States by the rat flea <i>Xenopsylla cheopis</i>).....	2481
Typhus fever. (The rat flea <i>Xenopsylla cheopis</i> in experimental transmission).....	1869
Typhus fever. Typhus virus in feces of infected fleas (<i>Xenopsylla cheopis</i>) and duration of infectivity of fleas.....	3103

Dysentery: (See Summaries—Disease cases reported monthly by States.)

E

Earle, W. R.:

A technique for adjustment of pH of hanging drop tissue cultures.....	1998
Technique for adjustment of the pH of tissue cultures planted in carrel flasks.....	2668

Ecuador: Guayaquil—Deaths—1930.....	2026
-------------------------------------	------

Employees' mutual benefit associations, a survey of the work of—Brundage.....	2102
---	------

Employees:

Sickness among male industrial, in the first quarter of 1931—Brundage.....	1799
--	------

Sickness among male industrial, in the second quarter of 1931—Brundage.....	2499
---	------

European conference on rural hygiene, held at Geneva, Switzerland, June 29—July 7, 1931.....	2671
--	------

Evans, Alice C.: New subspecies, <i>radicans</i> , of <i>alcaligenes faecalis</i>	Page 1676
The effect of hemolytic streptococci and their products on leucocytes.....	2539

F

Ferguson, G. H.: Sleeping-car parking and sanitation at a large convention.....	2229
Firestone, F.: Outbreaks of food poisoning apparently due to <i>B. enteritidis</i> , <i>B. paratyphosus B</i> (aertrycke type), and <i>B. paratyphosus A</i> , respectively.....	1565
Fleas, typhus virus in feces of infected, and duration of infectivity of—Ceder, Dyer, Rumreich, and Badger.....	3103
Florida: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.) Food poisoning, outbreaks of, apparently due to <i>B. enteritidis</i> , <i>B. paratyphosus B</i> (aertrycke type), and <i>B. paratyphosus A</i> , respectively—Geiger, Nelson, Gray, and Firestone.....	1565
Foreign reports: Communicable diseases.....	1601,
1658, 1713, 1786, 1856, 1896, 1967, 2027, 2078, 2145, 2208, 2276, 2341, 2399, 2465, 2523, 2585, 2647, 2693, 2759, 2823, 2882, 2958, 3008, 3087, 3140.	
France: Paris—Permanent committee of the International Office of Public Hygiene, May, 1931.....	2726
Francis, Edward: Agglutinin absorption in undulant fever (Brucellosis) ..	2416
French West Africa: Ivory Coast—Yellow fever.....	1965
Upper Volta—Yellow fever.....	1965
Frisbie, W. S.: Coordination in the sanitary control of bottled mineral waters.....	1873
Fumigants used for the destruction of cockroaches, experiments with certain—Ridlon.....	1623
Fumigation of vessels at Dairen and Port Arthur, Manchuria.....	2144
Fumigation on ships, effect of, on cockroaches—Williams.....	1680
Fumigation of loaded ships—Williams.....	1823
Fumigation of vessels, notes on. (Preliminary inspection, how rats escape, increased periods of exposure)—Williams.....	2973
Fumigation, report on some tests of the use of a new cyanogen product in ship—Williams.....	2048
Fumigation, ship, some aspects of—Ridlon.....	1572

G

Geiger, J. C.: Outbreaks of food poisoning apparently due to <i>B. enteritidis</i> , <i>B. paratyphosus B</i> (aertrycke type), and <i>B. paratyphosus A</i> , respectively.....	1565
Georgia: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.) German measles. (See Summaries—Disease cases reported monthly by States.)	
Gold Coast: Wale Wale—Yellow fever.....	1965
Government in public-health education, functions and limitations of—McLaughlin.....	2300
Gray, J. P.: Outbreaks of food poisoning apparently due to <i>B. enteritidis</i> , <i>B. paratyphosus B</i> (aertrycke type), and <i>B. paratyphosus A</i> , respectively.....	1565
Great Britain: England and Wales— Communicable diseases— Quarter ended July 4, 1931.....	2397
Quarter ended October 3, 1931.....	3139
Vital statistics— April-June, 1931.....	2396
July-September, 1931.....	3138
Scotland—Vital statistics— Quarter ended March 31, 1931.....	1785
Quarter ended June 30, 1931.....	2274
Quarter ended September 30, 1931.....	3139

	Page
Griffitts, J. J.: Mosquitoes transported by airplanes. (Staining method used in their importation).....	2775
Griffitts, T. H. D.: Anopheles atropos D. and K.—A new potential carrier of malaria organisms.....	3107
Mosquitoes transported by airplanes. (Staining method used in determining their importation).....	2775

H

Hasseltine, H. E.: Outbreak of undulant fever traced to infected milk supply.....	2291
Hawaii Territory: Kula district—Maui Island—Plague (human).....	1966
Health department and the medical profession—McLaughlin.....	2041
Health officers: City, 1931, directory of.....	2921
County, 1931, directory of.....	3051
State and insular, 1931, directory of.....	2899
Health of the school child.....	2605
Health program, rural, some essential considerations in connection with—Draper.....	1617
Hedrich, A. W.: Movements of epidemic meningitis, 1915-1930.....	2709
Hemolytic streptococci and their products on leucocytes, effect of—Evans.....	2539
Hiscock, Ira V.: A study of illness among grade-school children.....	1801
Holsendorf, B. E.: An important source of original rat infestation on newly constructed vessels.....	1943
Hookworm disease. (See Summaries—Disease cases reported monthly by States.)	
Hydrocyanic acid sprayer, air jet—Williams.....	1755

I

Idaho: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.)	
Illinois: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.)	
Illness among grade-school children, a study of—Wilson, Hiscock, Watkins, Case, and Rice.....	1801
Indiana: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.)	
Infantile paralysis. (See Poliomyelitis.)	
Infant mortality, birth and death figures, provisional, birth registration area, 1930.....	2373
Infection, double, by organisms of the Brucella group—Jordan.....	2437
Influence of temperature on the infecting power of Aedes aegypti containing the yellow fever virus.....	2739
Influenza: (See also Summaries—Disease cases reported monthly by States.) City reports for week ended—	
June 13, 1931.....	1590
June 20, 1931.....	1647
June 27, 1931.....	1702
July 4, 1931.....	1774
July 11, 1931.....	1844
July 18, 1931.....	1885
July 25, 1931.....	1955
August 1, 1931.....	2017
August 8, 1931.....	2067
August 15, 1931.....	2134
August 22, 1931.....	2195
August 29, 1931.....	2264
September 5, 1931.....	2329
September 12, 1931.....	2385
September 19, 1931.....	2453
September 26, 1931.....	2511

Influenza—Continued.

City reports for week ended—Continued.	Page
October 3, 1931.....	2572
October 10, 1931.....	2635
October 17, 1931.....	2681
October 24, 1931.....	2748
October 31, 1931.....	2812
November 7, 1931.....	2870
November 14, 1931.....	2946
November 21, 1931.....	2996
November 28, 1931.....	3075
December 5, 1931.....	3127
Current weekly State reports.....	1585,
1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129, 2191,	2259,
2324, 2380, 2447, 2506, 2566, 2629, 2675, 2744, 2806, 2866,	2940,
2989, 3067, 3121.	
Samoa.....	2398
Influenza and pneumonia morbidity and mortality, age and sex incidence of, in the epidemic of 1928-29 with comparative data for the epidemic of 1918-19—Collins.....	1909
Insane, admissions to hospitals for the:	
July, 1929.....	3073
August, 1929.....	3124
International Office of Public Hygiene, permanent committee, Paris, May, 1931.....	2726
Intestinal parasites of 73 boys in the National Training School for Boys, Washington, D. C., microscopic examination for.....	2980
Iowa:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Iraq:	
Basra—Cholera.....	1966
Cholera.....	3007

J

Jamaica: Communicable diseases—Four weeks ended—	
June 20, 1931.....	1785
July 18, 1931.....	2206
August 15, 1931.....	2274
September 12, 1931.....	2464
October 10, 1931.....	2692
Johnson, J. M.: Catalytic action of copper in the oxidation of crystalline glutathione.....	2234
Jordan, Carl F.: Double infection by organisms of the brucella group (report of a case).....	2437

K

Kansas:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Kentucky: (See Name of disease—Current weekly State reports.)	
Knight, I. W.: Outbreak of undulant fever traced to infected milk supply..	2291

L

Latvia: Communicable diseases—	
January-June, 1931.....	2340
July, 1931.....	2522
August, 1931.....	2822
September, 1931.....	2880
Lead poisoning. (See Summaries—Disease cases reported monthly by States.)	
Leathers, W. S.: Need for continued study in public-health work.....	1727
Leprosy:	
(See also Summaries—Disease cases reported monthly by States.)	
A study of white blood cells and their relation to clinical progress— Badger.....	2782

Lethargic encephalitis:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—

Page

June 13, 1931.....	1595
June 20, 1931.....	1652
June 27, 1931.....	1707
July 4, 1931.....	1780
July 11, 1931.....	1849
July 18, 1931.....	1888
July 25, 1931.....	1961
August 1, 1931.....	2022
August 8, 1931.....	2073
August 15, 1931.....	2139
August 22, 1931.....	2201
August 29, 1931.....	2269
September 5, 1931.....	2335
September 12, 1931.....	2391
September 19, 1931.....	2459
September 26, 1931.....	2517
October 3, 1931.....	2578
October 10, 1931.....	2641
October 17, 1931.....	2687
October 24, 1931.....	2753
October 31, 1931.....	2817
November 7, 1931.....	2876
November 14, 1931.....	2952
November 21, 1931.....	3002
November 28, 1931.....	3080
December 5, 1931.....	3133

Leucocytes, effect of hemolytic streptococci and their products on—Evans. 2539

Lillie, R. D.:

Pathology of the eastern type of Rocky Mountain spotted fever.... 2840

Typhus fever. (The experimental transmission of endemic typhus of the United States by the rat flea *Xenopsylla cheopis*)..... 2481

Long, John D.: Cooperative campaign for the eradication of plague in Peru (final report)..... 2161

Louisiana:

(See also Name of disease—Current weekly State reports.)

(See also Summaries—Disease cases reported monthly by States.)

New Orleans—Typhus fever patient removed from vessel "Atenas" at. 2384

Mc

McLaughlin, Allan J.:

The functions and limitations of Government in public-health education..... 2300

The medical profession and the health department..... 2041

M

Maine:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Malaria:

(See also Summaries—Disease cases reported monthly by States.)

Mexico—Monterrey..... 3007

Malaria organisms, a new potential carrier of—Anopheles atropos D. and K.—Mayne and Griffiths..... 3107

Maryland:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Massachusetts:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Mayne, Bruce: Anopheles atropos D. and K.—A new potential carrier of malaria organisms..... 3107

Meaker, Verna Thornhill: Dental decay and corrections among school children of different ages..... 2608

Measles:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—	Page
June 13, 1931	1590
June 20, 1931	1647
June 27, 1931	1702
July 4, 1931	1774
July 11, 1931	1844
July 18, 1931	1885
July 25, 1931	1955
August 1, 1931	2017
August 8, 1931	2067
August 15, 1931	2134
August 22, 1931	2195
August 29, 1931	2264
September 5, 1931	2329
September 12, 1931	2385
September 19, 1931	2453
September 26, 1931	2511
October 3, 1931	2572
October 10, 1931	2635
October 17, 1931	2681
October 24, 1931	2748
October 31, 1931	2812
November 7, 1931	2870
November 14, 1931	2946
November 21, 1931	2996
November 28, 1931	3075
December 5, 1931	3127
Current weekly State reports	1585,
1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129, 2191, 2259, 2324,	
2380, 2447, 2506, 2566, 2629, 2675, 2744, 2806, 2866, 2940, 2989, 3067,	
3121.	

Medical profession and the health department—McLaughlin 2041

Meningitis, movements of epidemic, 1915-1930—Hedrich 2709

Meningococcus meningitis:

(See also Summaries—Disease cases reported monthly by States.)

China—Shanghai 1711

City reports for week ended—

June 13, 1931	1595
June 20, 1931	1652
June 27, 1931	1707
July 4, 1931	1780
July 11, 1931	1849
July 18, 1931	1890
July 25, 1931	1961
August 1, 1931	2022
August 8, 1931	2073
August 15, 1931	2139
August 22, 1931	2201
August 29, 1931	2269
September 5, 1931	2335
September 12, 1931	2391
September 19, 1931	2459
September 26, 1931	2517
October 3, 1931	2578
October 10, 1931	2641
October 17, 1931	2687
October 24, 1931	2753
October 31, 1931	2817
November 7, 1931	2876
November 14, 1931	2952
November 21, 1931	3002
November 28, 1931	3080
December 5, 1931	3133

Current weekly State reports 1585,

1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129, 2191, 2259,
2324, 2380, 2447, 2506, 2566, 2629, 2675, 2744, 2806, 2866, 2940,
2989, 3067, 3121.

Meningococcus meningitis—Continued.	Page
On steamship "President Wilson" at San Francisco from Honolulu, October 6, 1931.....	2521
Metropolitan Life Insurance Co.—Death rates in a group of insured per- sons—	
May, 1931.....	1762
June, 1931.....	2059
July, 1931.....	2253
August, 1931.....	2623
September, 1931.....	2860
Mexico:	
Monterrey—Malaria.....	3007
Tampico—Communicable diseases—	
June, 1931.....	1854
July, 1931.....	2206
August, 1931.....	2340
September, 1931.....	2692
October, 1931.....	2881
Vera Cruz—Deaths—June 1-28, 1931.....	1785
Michigan:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Mineral waters, coordination in the sanitary control of bottled—Frisbie..	1873
Minnesota:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Mississippi:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Missouri:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Mitchell, A. Graeme: Scarlet-fever streptococcus antitoxin in the treat- ment of scarlet fever.....	3023
Montana:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Mortality. (See Deaths.)	
Mortality figures, provisional birth, death and infant, birth registration area, 1930.....	2373
Mortality statistics, comparative current State.....	2120, 3059
Mosquito infestation, inspection of ships for determination of—Tanner..	2306
Mosquitoes transported by airplanes (staining method used in importa- tion)—Griffitts and Griffitts.....	2775
Mountin, Joseph W.: Public health service in Knox County, Tenn.....	1981
Mouse, use of the white, in research on yellow fever.....	2366
Mumps:	
(See also Summaries—Disease cases reported monthly by States.)	
City reports for week ended—	
June 13, 1931.....	1590
June 20, 1931.....	1647
June 27, 1931.....	1702
July 4, 1931.....	1774
July 11, 1931.....	1844
July 18, 1931.....	1885
July 25, 1931.....	1955
August 1, 1931.....	2017
August 8, 1931.....	2067
August 15, 1931.....	2134
August 22, 1931.....	2195
August 29, 1931.....	2264
September 5, 1931.....	2329
September 12, 1931.....	2385
September 19, 1931.....	2453
September 26, 1931.....	2511
October 3, 1931.....	2572
October 10, 1931.....	2635

Mumps—Continued.

City reports for week ended—Continued.

	Page
October 17, 1931.....	2681
October 24, 1931.....	2748
October 31, 1931.....	2812
November 7, 1931.....	2870
November 14, 1931.....	2946
November 21, 1931.....	2996
November 28, 1931.....	3075
December 5, 1931.....	3127

N

Nebraska:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Nelson, Margaret: Outbreaks of food poisoning due to *B. enteritidis*, *B. paratyphosus* B (aertrycke type), and *B. paratyphosus* A, respectively.. 1565

New Hampshire:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

New Jersey:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

New Mexico:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

New York:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

New York State regulation against poisonous substances for polishing kitchenware or silverware..... 1945

North Carolina:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

North Dakota:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Notifiable diseases. (See Communicable diseases.)

O

Ohio:

(See also Name of disease—Current weekly State reports.)

(See also Summaries—Disease cases reported monthly by States.)

Cleveland—Typhoid fever outbreak at..... 2383

Oklahoma:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Oregon:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

P

Panama Canal Zone: Communicable diseases—

May, 1931.....	1855
June, 1931.....	1966
July, 1931.....	2275
August, 1931.....	2583
September, 1931.....	2757
October, 1931.....	3086

Paratyphoid fever. (See Summaries—Disease cases reported monthly by States.)

Parking, sleeping car, and sanitation at a large convention—Ferguson... 2229

Pathology of the eastern type of Rocky Mountain spotted fever—Lillie.. 2840

Pellagra:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—

Page

June 13, 1931.....	1595
June 20, 1931.....	1652
June 27, 1931.....	1707
July 4, 1931.....	1780
July 11, 1931.....	1849
July 18, 1931.....	1890
July 25, 1931.....	1961
August 1, 1931.....	2022
August 8, 1931.....	2073
August 15, 1931.....	2139
August 22, 1931.....	2201
August 29, 1931.....	2269
September 5, 1931.....	2335
September 12, 1931.....	2391
September 19, 1931.....	2459
September 26, 1931.....	2517
October 3, 1931.....	2578
October 10, 1931.....	2641
October 17, 1931.....	2687
October 24, 1931.....	2753
October 31, 1931.....	2817
November 7, 1931.....	2876
November 14, 1931.....	2952
November 21, 1931.....	3002
November 28, 1931.....	3080
December 5, 1931.....	3133

Pellagra in the United States, a note on the history of—Wheeler..... 2223

Pellagra-preventive value of canned spinach, canned turnip greens, mature onions, and canned green beans—Wheeler..... 2663

Pennsylvania:

(See also Name of disease—Current weekly State reports.)

(See also Summaries—Disease cases reported monthly by States.)

West Chester—Outbreak of typhoid fever at the State Teachers College..... 2633

Permanent committee of the International Office of Public Hygiene, Paris, May, 1931..... 2726

Persia: Cholera, measures against..... 2692

Peru, cooperative campaign for the eradication of plague in—Long..... 2161

Philip, Cornelius B.: Occurrence of a colony of the tick parasite *Huntellus hookeri* Howard in West Africa..... 2168

Physical examination as an instrument of research—Britten..... 1671

Plague (human):

Argentina—San Juan Province..... 1894

China—

Changchow..... 1894

Chiobe..... 1894

Shansi Province..... 2646, 2821

Shensi Province..... 2821

Foreign reports..... 1603,

1660, 1715, 1788, 1858, 1898, 1970, 2029, 2080, 2147, 2210,

2278, 2344, 2402, 2468, 2526, 2588, 2650, 2696, 2762, 2826,

2885, 2961, 3011, 3090, 3143.

Hawaii Territory—Kula district—Maui Island..... 1966

Plague (rodent): California—San Benito County..... 1954, 2133

Plague in Peru, cooperative campaign for the eradication of—Long..... 2161

Pneumonia (all forms): City reports for week ended—

June 13, 1931.....	1590
June 20, 1931.....	1647
June 27, 1931.....	1702
July 4, 1931.....	1774
July 11, 1931.....	1844
July 18, 1931.....	1885
July 25, 1931.....	1955
August 1, 1931.....	2017
August 8, 1931.....	2067
August 15, 1931.....	2134

Pneumonia (all forms): City reports for week ended—Continued.	Page
August 22, 1931.....	2195
August 29, 1931.....	2264
September 5, 1931.....	2329
September 12, 1931.....	2385
September 19, 1931.....	2453
September 26, 1931.....	2511
October 3, 1931.....	2572
October 10, 1931.....	2635
October 17, 1931.....	2681
October 24, 1931.....	2748
October 31, 1931.....	2812
November 7, 1931.....	2870
November 14, 1931.....	2946
November 21, 1931.....	2996
November 28, 1931.....	3075
December 5, 1931.....	3127
Pneumonia and influenza morbidity and mortality, age and sex incidence of, in the epidemic of 1928-29 with comparative data for the epidemic of 1918-19—Collins.....	1909
Poisonous substances for polishing kitchenware or silverware, New York State regulation against.....	1945
Poliomyelitis:	
(See also Summaries—Disease cases reported monthly by States.)	
City reports for week ended—	
June 13, 1931.....	1595
June 20, 1931.....	1652
June 27, 1931.....	1707
July 4, 1931.....	1780
July 11, 1931.....	1849
July 18, 1931.....	1888
July 25, 1931.....	1961
August 1, 1931.....	2022
August 8, 1931.....	2073
August 15, 1931.....	2139
August 22, 1931.....	2201
August 29, 1931.....	2269
September 5, 1931.....	2335
September 12, 1931.....	2391
September 19, 1931.....	2459
September 26, 1931.....	2517
October 3, 1931.....	2578
October 10, 1931.....	2641
October 17, 1931.....	2687
October 24, 1931.....	2753
October 31, 1931.....	2817
November 7, 1931.....	2876
November 14, 1931.....	2952
November 21, 1931.....	3002
November 28, 1931.....	3080
December 5, 1931.....	3133
Current weekly State reports.....	1586,
1644, 1698, 1769, 1841, 1881, 1951, 2014, 2064, 2130, 2192, 2260,	
2325, 2381, 2448, 2507, 2567, 2630, 2676, 2745, 2807, 2867, 2941,	
2990, 3068, 3122.	
Porto Rico: San Juan—Communicable diseases—Four weeks ended—	
June 20, 1931.....	1855
July 18, 1931.....	2206
August 15, 1931.....	2398
September 12, 1931.....	2522
October 10, 1931.....	2758
November 7, 1931.....	2881
Publications, Public Health Service. List of publications issued during the period January-June, 1931.....	1636
Public health—Court decisions relating to.....	1582,
1694, 1763, 1836, 1876, 1946, 2008, 2060, 2125, 2187, 2320, 2376,	
2443, 2502, 2562, 2625, 2740, 2802, 2984, 3064, 3117.	

	Page
Public health education, functions and limitations of government in— McLaughlin.....	2300
Public health service in Knox county, Tenn.—Mountin.....	1981
Public Health Service publications. List of publications issued during the period January-June, 1931.....	1636
Public health work, need for continued study in—Leathers.....	1727
Puerperal fever. (See Summaries—Disease cases reported monthly by States.)	
R	
Rabies in animals. (See Summaries—Disease cases reported monthly by States.)	
Rabies in man. (See Summaries—Disease cases reported monthly by States.)	
Radicans, a new subspecies, of <i>Alcaligenes faecalis</i> —Evans.....	1676
Rat flea (<i>Xenopsylla cheopis</i>) experimental transmission of endemic typhus fever of the United States, by the—Dyer, Ceder, Rumreich, and Badger.....	2415
Rat-flea survey of the port of St. Thomas, Virgin Islands—Carnes.....	2558
Rat infestation on newly constructed vessels, an important source of original—Holsendorf.....	1943
Rat population of Diesel motor boats.....	2371
Reciprocal notifications of communicable disease cases—United States— April, 1931.....	2194
May, 1931.....	2262
June, 1931.....	2328
July, 1931.....	2384
August, 1931.....	2452
September, 1931.....	2633
October, 1931.....	2945
Regulation, New York State, against poisonous substances for polishing kitchenware or silverware.....	1945
Rhode Island: (See Name of disease—Current weekly State reports.) (See Summaries—Disease cases reported monthly by States.)	
Rice, John L: A study of illness among grade-school children.....	1801
Ridlon, J. R.: Experiments with certain fumigants used for the destruction of cock- roaches.....	1623
Some aspects of ship fumigation.....	1572
Ringworm. (See Summaries—Disease cases reported monthly by States.)	
Rocky Mountain spotted fever: (See also Summaries—Disease cases reported monthly by States.) Pathology of the eastern type of—Lillie.....	2840
Rosenthal, Sanford M.: Catalytic action of copper in the oxidation of crystalline glutathione.....	2234
Rumreich, A.: Experimental transmission of endemic typhus fever of the United States by the rat flea (<i>Xenopsylla cheopis</i>).....	2415
Typhus fever. (The experimental transmission of endemic typhus fever of the United States by the rat flea (<i>Xenopsylla cheopis</i>).....	2481
Typhus fever. (The rat flea <i>Xenopsylla cheopis</i> in experimental transmission).....	1869
Typhus fever. Typhus virus in feces of infected fleas (<i>Xenopsylla cheopis</i>) and duration of infectivity of fleas.....	3103
Rural health program, some essential considerations in connection with— Draper.....	1617
Rural health service in the United States, 1927-1931.....	2173
Rural hygiene, European conference on, held at Geneva, Switzerland, June 29-July 7, 1931.....	2671
S	
Samoa: Influenza.....	2398
Sanitary control of bottled mineral waters, coordination in the—Frisbie..	1873
Sanitation and sleeping car parking at a large convention—Ferguson.....	2229
Scabies. (See Summaries—Disease cases reported monthly by States.)	

Scarlet fever:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—

	Page
June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750
October 31, 1931.....	2815
November 7, 1931.....	2873
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130

Current weekly State reports..... 1586, 1644,

1698, 1769, 1841, 1881, 1951, 2014, 2064, 2130, 2192, 2260, 2325, 2381,

2448, 2507, 2567, 2630, 2676, 2745, 2807, 2867, 2941, 2990, 3068, 3122

Scarlet fever streptococcus antitoxin in the treatment of scarlet fever—

Veldee, Stevenson, and Mitchell..... 3023

Schwartz, Louis: Dermatitis venenata due to contact with Brazilian walnut wood..... 1938

Septic sore throat. (See Summaries—Disease cases reported monthly by States.)

Ship fumigation, some aspects of—Ridlon..... 1572

Ships, fumigation of loaded—Williams..... 1823

Ships, inspection of, for determination of mosquito infestation—Tanner..... 2306

Sickness among male industrial employees in the first quarter of 1931—

Brundage..... 1799

Sickness among male industrial employees in the second quarter of 1931—

Brundage..... 2499

Smallpox:

(See also Summaries—Disease cases reported monthly by States.)

City reports for week ended—

June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750

Smallpox—Continued.

City reports for week ended—Continued.

	Page
October 31, 1931.....	2815
November 7, 1931.....	2873
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130

Current weekly State reports.

1644, 1698, 1769, 1841, 1881, 1951, 2014, 2064, 2130, 2192, 2260, 2325, 2381, 2448, 2507, 2567, 2630, 2676, 2745, 2807, 2867, 2941, 2990, 3068, 3122.	1586,
---	-------

Foreign reports.

1663, 1718, 1791, 1861, 1901, 1973, 2033, 2083, 2150, 2213, 2282, 2347, 2405, 2471, 2530, 2591, 2653, 2700, 2765, 2829, 2889, 2964, 3014, 3094, 3147.	1606,
---	-------

South America: Yellow fever..... 1895

South Carolina:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

South Dakota:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Spencer, R. R.: Expansion of investigations on tick-borne diseases by the

United States Public Health Service..... 2097

Sprayer, air jet hydrocyanic acid—Williams..... 1755

State and insular health authorities, 1931, directory of..... 2899

Statistics, comparative current State mortality..... 2120, 3059

Stevenson, F. E.: Scarlet-fever streptococcus antitoxin in the treatment of scarlet fever..... 3023

Stoughton, Amanda L.: Dental decay and corrections among school children of different ages..... 2608

Summaries:

Communicable diseases—

Cases and case rates per 100,000 population—Reported by State health officers—

February, 1931.....	1771, 1772
March, 1931.....	2451, 2452
April, 1931.....	2509, 2510
May, 1931.....	2569, 2570
June, 1931.....	2679, 2680
July, 1931.....	2810, 2811
August, 1931.....	2993, 2994
September, 1931.....	3071, 3072

Case rates per 100,000 population for 98 cities—

May 10–June 13, 1931.....	1597
May 17–June 20, 1931.....	1654
May 24–June 27, 1931.....	1709
May 31–July 4, 1931.....	1782
June 7–July 11, 1931.....	1851
June 14–July 18, 1931.....	1892
June 21–July 25, 1931.....	1962
June 28–August 1, 1931.....	2024
July 5–August 8, 1931.....	2075
July 12–August 15, 1931.....	2141
July 19–August 22, 1931.....	2202
July 26–August 29, 1931.....	2271
August 2–September 5, 1931.....	2337
August 9–September 12, 1931.....	2393
August 16–September 19, 1931.....	2461
August 23–September 26, 1931.....	2519
August 30–October 3, 1931.....	2580
September 6–October 10, 1931.....	2643
September 13–October 17, 1931.....	2688
September 20–October 24, 1931.....	2755
September 27–October 31, 1931.....	2819
October 4–November 7, 1931.....	2878
October 11–November 14, 1931.....	2954
October 18–November 21, 1931.....	3004

Summaries—Continued.

Communicable diseases—Continued.

Case rates per 100,000 population for 98 cities—Continued.	Page
October 25–November 28, 1931.....	3082
November 1–December 5, 1931.....	3135
Disease cases reported monthly by States.....	1587,
1645, 1699, 1770, 1842, 1882, 1952, 2015, 2065, 2131, 2193, 2261,	2261,
2326, 2382, 2449, 2508, 2568, 2631, 2677, 2746, 2808, 2868, 2942,	2942,
2991, 3069, 3123.	
General current, and weekly reports from cities—Week ended—	
June 13, 1931.....	1598
June 20, 1931.....	1646
June 27, 1931.....	1701
July 4, 1931.....	1773
July 11, 1931.....	1843
July 18, 1931.....	1884
July 25, 1931.....	1954
August 1, 1931.....	2016
August 8, 1931.....	2067
August 15, 1931.....	2133
August 22, 1931.....	2194
August 29, 1931.....	2263
September 5, 1931.....	2328
September 12, 1931.....	2384
September 19, 1931.....	2453
September 26, 1931.....	2510
October 3, 1931.....	2571
October 10, 1931.....	2633
October 17, 1931.....	2680
October 24, 1931.....	2747
October 31, 1931.....	2811
November 7, 1931.....	2870
November 14, 1931.....	2945
November 21, 1931.....	2995
November 28, 1931.....	3074
December 5, 1931.....	3126

Switzerland:

Geneva—European conference on rural hygiene, held at, June 29–July 7, 1931.....	2671
Tuberculosis, deaths from, 1911–1920, 1921–1930.....	3139

T

Tanner, W. F.: Inspection of ships for determination of mosquito infestation.....	2306
Tasmania: Vital statistics, 1930.....	2583
Tennessee:	
(See also Name of disease—Current weekly State reports.)	
(See also Summaries—Disease cases reported monthly by States.)	
Knox County, public health service in—Mountain.....	1981
Tetanus. (See Summaries—Disease cases reported monthly by States.)	
Texas:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Tick-borne diseases, expansion of investigations on, by the United States Public Health Service—Spencer.....	2097
Tick parasite <i>Hunterellus hookeri</i> Howard in West Africa, occurrence of a colony of—Philip.....	2168
Tissue cultures planted in carrel flasks, technique for adjustment of the pH of—Earle.....	2668
Trachoma. (See Summaries—Disease cases reported monthly by States.)	
Trinidad: Port of Spain—Vital statistics—	
May, 1931, 1930.....	1712
June, 1931, 1930.....	2207
July, 1931, 1930.....	2340
August, 1931, 1930.....	2464
September, 1931, 1930.....	2822
October, 1931, 1930.....	3086
Tropical medicine, special course in clinical.....	1640

Tuberculosis (all forms): City reports for week ended—	Page
June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750
October 31, 1931.....	2815
November 7, 1931.....	2873
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130
Tuberculosis: Switzerland (deaths in), 1911-1920, 1921-1930.....	3139
Tularaemia. (See Summaries—Disease cases reported monthly by States.)	
Typhoid fever:	
(See also Summaries—Disease cases reported monthly by States.)	
City reports for week ended—	
June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750
October 31, 1931.....	2815
November 7, 1931.....	2873
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130
Current weekly State reports.....	1586,
1644, 1698, 1769, 1841, 1881, 1951, 2014, 2064, 2130, 2192, 2260,	
2325, 2381, 2448, 2507, 2567, 2630, 2676, 2745, 2807, 2867, 2941,	
2990, 3068, 3122.....	
Ohio—Cleveland.....	2383
Typhoid fever outbreak at State Teachers College, West Chester, Pa.	2633

Typhus fever:

(See also City reports.)

(See also Summaries—Disease cases reported monthly by States.)

Foreign reports	Page
1668, 1723, 1796, 1865, 1905, 1978, 2038, 2088, 2155, 2218, 2286, 2352, 2410, 2476, 2534, 2596, 2658, 2704, 2770, 2833, 2893, 2968, 3018, 3098, 3151.	1611,

Typhus fever, experimental transmission of endemic, of the United States by the rat flea *Xenopsylla cheopis*—Dyer, Ceder, Rumreich, and Badger

2415

Typhus fever; experimental transmission of endemic of the United States by the rat flea *Xenopsylla cheopis*—Dyer, Ceder, Lillie, Rumreich, and Badger

2481

Typhus fever. (The rat flea *Xenopsylla cheopis* in experimental transmission)—Dyer, Ceder, Rumreich, and Badger

1869

Typhus fever: Typhus virus in feces of infected fleas (*Xenopsylla cheopis*) and duration of infectivity of fleas—Ceder, Dyer, Rumreich, and Badger

3103

Typhus fever patient removed from vessel "Atenas" at New Orleans, La.

2384

U

Undulant fever:

(See also Summaries—Disease cases reported monthly by States.)

Agglutinin absorption in (Brucellosis)—Francis

2416

Outbreak of, traced to infected milk supply—Hasseltine and Knight

2291

United States:

Communicable diseases—

Cases and case rates per 100,000 population—Reported by State health officers—

February, 1931	1771, 1772
March, 1931	2451, 2452
April, 1931	2509, 2510
May, 1931	2569, 2570
June, 1931	2679, 2680
July, 1931	2810, 2811
August, 1931	2993, 2994
September, 1931	3071, 3072

Case rates per 100,000 population for 98 cities—

May 10–June 13, 1931	1597
May 17–June 20, 1931	1654
May 24–June 27, 1931	1709
May 31–July 4, 1931	1782
June 7–July 11, 1931	1851
June 14–July 18, 1931	1892
June 21–July 25, 1931	1962
June 28–August 1, 1931	2024
July 5–August 8, 1931	2075
July 12–August 15, 1931	2141
July 19–August 22, 1931	2202
July 26–August 29, 1931	2271
August 2–September 5, 1931	2337
August 9–September 12, 1931	2393
August 16–September 19, 1931	2461
August 23–September 26, 1931	2519
August 30–October 3, 1931	2580
September 6–October 10, 1931	2643
September 13–October 17, 1931	2688
September 20–October 24, 1931	2755
September 27–October 31, 1931	2819
October 4–November 7, 1931	2878
October 11–November 14, 1931	2954
October 18–November 21, 1931	3004
October 25–November 28, 1931	3082
November 1–December 5, 1931	3135

Current prevalence of 1615, 1871, 2093, 2357, 2601, 2837, 3115

Current weekly State reports 1585,

1643, 1697, 1768, 1840, 1880, 1950, 2013, 2063, 2129, 2191, 2259, 2324, 2380, 2447, 2506, 2566, 2629, 2675, 2744, 2806, 2866, 2940, 2989, 3067, 3121.

United States—Continued.

Communicable diseases—Continued.

Deaths—Large cities—Week ended—

	Page
June 13, 1931.....	1582
June 20, 1931.....	1640
June 27, 1931.....	1694
July 4, 1931.....	1765
July 11, 1931.....	1837
July 18, 1931.....	1877
July 25, 1931.....	1947
August 1, 1931.....	2010
August 8, 1931.....	2060
August 15, 1931.....	2126
August 22, 1931.....	2188
August 29, 1931.....	2256
September 5, 1931.....	2321
September 12, 1931.....	2377
September 19, 1931.....	2444
September 26, 1931.....	2503
October 3, 1931.....	2563
October 10, 1931.....	2626
October 17, 1931.....	2672
October 24, 1931.....	2741
October 31, 1931.....	2803
November 7, 1931.....	2863
November 14, 1931.....	2937
November 21, 1931.....	2986
November 28, 1931.....	3064
December 5, 1931.....	3118

Reciprocal notifications of—

April, 1931.....	2194
May, 1931.....	2262
June, 1931.....	2328
July, 1931.....	2384
August, 1931.....	2452
September, 1931.....	2633
October, 1931.....	2945

Summaries—Disease cases reported monthly by States.....	1587,
1645, 1699, 1770, 1842, 1882, 1952, 2015, 2065, 2131, 2193, 2261,	
2326, 2382, 2449, 2508, 2568, 2631, 2677, 2746, 2808, 2868, 2942,	
2991, 3069, 3123.	

Note on the history of pellagra in—Wheeler..... 2223

United States, extent of rural health service in, 1927-1931..... 2173

Use of the white mouse in research on yellow fever..... 2366

Utah. (See Name of disease—Current weekly State reports.)

V

Veldee, M. V.: Scarlet-fever streptococcus antitoxin in the treatment of scarlet fever..... 3023

Vermont:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Vessels:

Newly constructed, an important source of original rat infestation on—Holsendorf..... 1943

Notes on the fumigation of. (Preliminary inspection, how rats escape, increased periods of exposure)—Williams..... 2973

Vincent's angina. (See Summaries—Disease cases reported monthly by States.)

Virginia:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Virgin Islands:

Communicable diseases—

May, 1931.....	1657
June, 1931.....	1855
July, 1931.....	2077

Virgin Islands—Continued.	
Communicable diseases—Continued.	Page
August, 1931.....	2398
September, 1931.....	2646
October, 1931.....	2957
St. Thomas—Rat-flea survey of—Carnes.....	2558
Vital statistics:	
Canada—Saskatchewan—1930.....	3006
China—Shansi Province—Year 1923.....	2582
Great Britain—	
England and Wales—	
April-June, 1931.....	2396
July-September, 1931.....	3138
Scotland—	
Quarter ended, March 31, 1931.....	1785
Quarter ended June 30, 1931.....	2274
Quarter ended September 30, 1931.....	3139
Tasmania, 1930.....	2583
Trinidad—Port of Spain—	
May, 1931, 1930.....	1712
June, 1931, 1930.....	2207
July, 1931, 1930.....	2340
August, 1931, 1930.....	2464
September, 1931, 1930.....	2822
October, 1931, 1930.....	3086
Voegtlin, Carl. Catalytic action of copper in the oxidation of crystalline glutathione.....	2234

W

Washington:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Watkins, J. H., A study of illness among grade-school children.....	1801
West Africa, occurrence of a colony of tick parasite <i>Hunterellus hookeri</i> Howard in—Philip.....	2168
West Virginia:	
(See Name of disease—Current weekly State reports.)	
(See Summaries—Disease cases reported monthly by States.)	
Wheeler, G. A.:	
Note on the history of pellagra in the United States.....	2223
The pellagra-preventive value of canned spinach, canned turnip greens, mature onions, and canned green beans.....	2663
Whooping cough:	
(See also Summaries—Disease cases reported monthly by States.)	
City reports for week ended—	
June 13, 1931.....	1593
June 20, 1931.....	1650
June 27, 1931.....	1705
July 4, 1931.....	1777
July 11, 1931.....	1847
July 18, 1931.....	1888
July 25, 1931.....	1958
August 1, 1931.....	2020
August 8, 1931.....	2070
August 15, 1931.....	2136
August 22, 1931.....	2198
August 29, 1931.....	2267
September 5, 1931.....	2332
September 12, 1931.....	2388
September 19, 1931.....	2456
September 26, 1931.....	2514
October 3, 1931.....	2575
October 10, 1931.....	2638
October 17, 1931.....	2684
October 24, 1931.....	2750
October 31, 1931.....	2815
November 7, 1931.....	2873

Whooping cough—Continued.

City reports for week ended—Continued.

	Page
November 14, 1931.....	2949
November 21, 1931.....	2999
November 28, 1931.....	3078
December 5, 1931.....	3130

Williams, C. L.:

Air jet hydrocyanic acid sprayer.....	1755
Effect of fumigation on cockroaches on ships.....	1680
Fumigation of loaded ships.....	1823
Notes on the fumigation of vessels. (Preliminary inspection, how rats escape, increased periods of exposure).....	2973
Report on some tests of the use of a new cyanogen product in ship fumigation.....	2048

Wilson, Charles C.: A study of illness among grade-school children.....

1801

Wisconsin:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Wynns, H. L.: Outbreaks of food poisoning apparently due to *B. enteritidis*, *B. paratyphosus B* (aertrycke type), and *B. paratyphosus A*, respectively.....

1565

Wyoming:

(See Name of disease—Current weekly State reports.)

(See Summaries—Disease cases reported monthly by States.)

Y

Yellow fever:

British Cameroons—Mamfe, May 28, 1931.....

1656

Foreign reports.....

1613,

1670, 1725, 1798, 1868, 1908, 1980, 2040, 2090, 2158, 2220, 2288,

2354, 2412, 2478, 2536, 2598, 2660, 2706, 2772, 2835, 2896, 2970,

3020, 3100, 3153.

French West Africa—

Ivory Coast.....

1965

Upper Volta.....

1965

Gold Coast—Wale Wale.....

1965

Present-day problems of—Cumming.....

2361

South America.....

1895

Use of the white mouse in research on.....

2366

Yellow fever virus, influence of temperature on the infecting power of

Aedes aegypti containing.....

2739

Yugoslavia: Communicable diseases—

May, 1931.....

1600

June, 1931.....

2207

July, 1931.....

2275

August, 1931.....

2398

September, 1931.....

2822

October, 1931.....

3007

